

International Ocean Discovery Program
Expedition 383 Preliminary Report
Dynamics of the Pacific Antarctic Circumpolar Current
(DYNAPACC)

20 May–20 July 2019

Frank Lamy, Gisela Winckler, Carlos A. Alvarez Zarikian, and the Expedition 383 Scientists

Publisher's notes

Core samples and the wider set of data from the science program covered in this report are under moratorium and accessible only to Science Party members until 18 July 2021.

This publication was prepared by the *JOIDES Resolution* Science Operator (JRSO) at Texas A&M University (TAMU) as an account of work performed under the International Ocean Discovery Program (IODP). This material is based upon work supported by the JRSO, which is a major facility funded by the National Science Foundation Cooperative Agreement number OCE1326927. Funding for IODP is provided by the following international partners:

National Science Foundation (NSF), United States
Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan
European Consortium for Ocean Research Drilling (ECORD)
Ministry of Science and Technology (MOST), People's Republic of China
Korea Institute of Geoscience and Mineral Resources (KIGAM)
Australia-New Zealand IODP Consortium (ANZIC)
Ministry of Earth Sciences (MoES), India
Coordination for Improvement of Higher Education Personnel (CAPES), Brazil

Portions of this work may have been published in whole or in part in other IODP documents or publications.

Disclaimer

The *JOIDES Resolution* Science Operator is supported by the National Science Foundation. Any opinions, findings, and conclusions or recommendations expressed in this material do not necessarily reflect the views of the National Science Foundation, the participating agencies, TAMU, or Texas A&M Research Foundation.

Copyright

Except where otherwise noted, this work is licensed under a Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>). Unrestricted use, distribution, and reproduction are permitted, provided the original author and source are credited.



Citation

Lamy, F., Winckler, G., Alvarez Zarikian, C.A., and the Expedition 383 Scientists, 2019. *Expedition 383 Preliminary Report: Dynamics of the Pacific Antarctic Circumpolar Current*. International Ocean Discovery Program.
<https://doi.org/10.14379/iodp.pr.383.2019>

ISSN

World Wide Web: 2372-9562

Expedition 383 participants

Expedition 383 scientists

Frank Lamy

Co-Chief Scientist

Department of Marine Geology
Alfred Wegener Institute Helmholtz-Center for Polar and
Marine Research
Germany
Frank.Lamy@awi.de

Gisela Winckler

Co-Chief Scientist

Lamont-Doherty Earth Observatory
Columbia University
USA
winckler@ldeo.columbia.edu

Carlos A. Alvarez Zarikian

Expedition Project Manager/Staff Scientist

International Ocean Discovery Program
Texas A&M University
USA
zarikian@iodp.tamu.edu

Helge W. Arz

Sedimentologist

Marine Geology Department
Leibniz Institute for Baltic Sea Research Warnemünde
Germany
helge.arz@io-warnemuende.de

Chandranath Basak

Inorganic Geochemist

Department of Earth Sciences
University of Delaware
USA
cbasak@udel.edu

Anieke Brombacher

Paleontologist (foraminifers)

School Ocean & Earth Science
National Oceanography Centre
United Kingdom
anieke.brombacher@soton.ac.uk

Oliver M. Esper

Paleontologist (diatoms/dinoflagellates)

Department of Marine Geology
Alfred Wegener Institute Helmholtz-Center for Polar and
Marine Research
Germany
oliver.esper@awi.de

Jesse R. Farmer

Sedimentologist/Core Description

Department of Geological Sciences
Princeton University
USA
jesse.farmer@princeton.edu

Julia Gottschalk

Sedimentologist
Lamont-Doherty Earth Observatory
Columbia University
USA
jgottsch@ldeo.columbia.edu

Lisa C. Herbert

Inorganic Geochemist

School of Marine and Atmospheric Sciences
Stony Brook University
USA
lisa.herbert@stonybrook.edu

Shinya Iwasaki

Sedimentologist

Research and Development Center for Global Change
JAMSTEC
Japan
iwasaki.shinya@jamstec.go.jp

Vera J. Lawson

Physical Properties Specialist/Petrophysics

Department of Earth and Planetary Sciences
Rutgers, The State University of New Jersey
USA
vera.lawson@rutgers.edu

Lester Lembke-Jene

Physical Properties Specialist/Petrophysics

Department of Geosciences
Alfred Wegener Institute Helmholtz-Center for Polar and
Marine Research
Germany
lester.lembke-jene@awi.de

Li Lo

Sedimentologist

Department of Geosciences
National Taiwan University
lilo115@ntu.edu.tw

Alexandre L. de Souza

Paleontologist (diatoms/radiolarians)

Department of Geology
Federal University of Rio de Janeiro
Brazil
alsouza.bio@gmail.com

Elisa Malinverno

Paleontologist (nannofossils)

Department of Earth and Environmental Sciences
University of Milano-Bicocca
Italy
elisa.malinverno@unimib.it

Elisabeth M.L. Michel**Physical Properties Specialist/Petrophysics**

Laboratoire des Sciences du Climat et de l'Environnement LSCE
 Institut Pierre Simon Laplace IPSL
 CNRS-CEA-UVSQ
 France
Elisabeth.michel@lsce.ipsl.fr

Jennifer Middleton**Stratigraphic Correlator**

Lamont-Doherty Earth Observatory
 Columbia University
 USA
jennym@ldeo.columbia.edu

Simone Moretti**Organic Geochemist**

Climate Geochemistry Department
 Max Planck Institute for Chemistry (MPIC)
 Germany
simone.moretti@mpic.de

Christopher M. Moy**Sedimentologist**

Geology Department
 University of Otago
 New Zealand
chris.moy@otago.ac.nz

Christina Ravelo**Stratigraphic Correlator**

Ocean Sciences Department
 University of California, Santa Cruz
 USA
acr@ucsc.edu

Christina Riesselman**Paleontologist (diatoms)**

Geology Department
 University of Otago
 New Zealand
christina.riesselman@otago.ac.nz

Mariem Saavedra Pellitero**Paleontologist (nannofossils)**

School of Geography, Earth and Environmental Sciences
 University of Birmingham
 United Kingdom
msaavedr@uni-bremen.de

Observer**Nicole Foucher McColl**

Universidad Nacional Andrés Bello, UNAB
 Chile
n.fouchermccoll@uandresbello.edu

Outreach**Sian H. Proctor****Outreach Officer**

Geoscience Department
 South Mountain Community College
 USA
sian.proctor@southmountaincc.edu

Inah Seo**Sedimentologist**

Global Ocean Research Center
 Korea Institute of Ocean Science and Technology (KIOST)
 Republic of Korea
inahseo@kiost.ac.kr

Raj K. Singh**Paleontologist (foraminifers)**

School of Earth, Ocean and Climate Sciences
 Indian Institute of Technology Bhubaneswar
 India
rksingh@iitbbs.ac.in

Rebecca A. Smith**Organic Geochemist**

Geosciences
 University of Massachusetts-Amherst
 USA
rasmith@geo.umass.edu

Joseph S. Stoner**Paleomagnetist**

College of Earth, Ocean and Atmospheric Sciences
 Oregon State University
 USA
jstoner@coas.oregonstate.edu

Igor M. Venancio P de Oliveira**Paleontologist (foraminifers)**

Center for Weather Forecasting and Climate Research (CPTEC)
 National Institute for Spatial Research (INPE-Brazil)
 Brazil
igor.venancio@inpe.br

Sui Wan**Sedimentologist**

South China Sea Institute of Oceanology
 Chinese Academy of Sciences
 China
wansui@scsio.ac.cn

Xiangyu Zhao**Paleomagnetist**

National Institute of Polar Research
 Geoscience Group
 Japan
zhao.xiangyu@nipr.ac.jp

Brian Cimagala**Film Producer**

Maylo Films
 USA
Bcimagala@gmail.com

Alejandro Wilkins
Director of Photography
Maylo Films
USA
alejandro.image@gmail.com

Siem Offshore AS officials

Jacob C. Robinson
Master of the Drilling Vessel

Mark Robinson
Drilling Supervisor

Technical support

Timothy Blaisdell
Application Developer

Nicolette Lawler
X-ray Laboratory

Susan Boehm
Thin Section Laboratory

Chang Liu
Chemistry Laboratory

Ty Cobb
X-ray Laboratory

Daniel Marone
Underway Geophysics Specialist

Aaron de Loach
Assistant Laboratory Officer

Aaron Mechler
Chemistry Laboratory

Ekant Desai
Publications Specialist

Mike Meiring
Electronics Specialist

Timothy Fulton
Imaging Specialist

Beth Novak
Paleomagnetism Laboratory

Clayton Furman
Logging Engineer

Elisabetta Olivo
Marine Laboratory Specialist (temporary)

Randy Gjesvold
Marine Instrumentation Specialist

Chieh Peng
Laboratory Officer

Kevin Grigar
Operations Superintendent

Ionela Samoila
Marine Laboratory Specialist (temporary)

Sandra Herrmann
Assistant Laboratory Officer

Catherine Smith
Curatorial Specialist

Michael Hodge
Marine Computer Specialist

Steven Thomas
Marine Computer Specialist

Jon Howell
Application Developer

Garrick Van Rensburg
Marine Instrumentation Specialist

Skyler Jordan
Marine Laboratory Specialist (temporary)

Abstract

The Antarctic Circumpolar Current (ACC) is the world's strongest zonal current system that connects all three major ocean basins of the global ocean and therefore integrates and responds to global climate variability. Its flow is largely driven by strong westerly winds and constricted to its narrowest extent in the Drake Passage. Transport of fresh and cold surface and intermediate water masses through the Drake Passage (cold-water route) strongly affects the Atlantic Meridional Overturning Circulation together with the inflow of Indian Ocean water masses (warm-water route). Both oceanographic corridors are critical for the South Atlantic contribution to Meridional Overturning Circulation changes. In contrast to the Atlantic and Indian sectors of the ACC, and with the exception of drill cores from the Antarctic continental margin and off New Zealand, the Pacific sector of the ACC lacks information on its Cenozoic paleoceanography from deep-sea drilling records. To advance our knowledge and understanding of Miocene to Holocene atmosphere-ocean-cryosphere dynamics in the Pacific and their implications for regional and global climate and atmospheric CO₂, International Ocean Discovery Program (IODP) Expedition 383 recovered sedimentary sequences at (1) three sites located in the central South Pacific (U1539, U1540, and U1541), (2) two sites at the Chile margin (U1542 and U1544), and (3) one site from the pelagic eastern South Pacific (U1543) close to the entrance to the Drake Passage. Because of persistently stormy conditions and the resulting bad weather avoidance, we were not successful in recovering the originally planned Proposed Site CSP-3A in the central South Pacific in the Polar Frontal Zone. The drilled sediments at Sites U1541 and U1543 reach back to the late Miocene, and those at Site U1540 reach back to the early Pliocene. High sedimentary rate Pleistocene sedimentary sequences were drilled both in the central South Pacific (Site U1539) and along the Chile margin. Taken together, the sites represent a depth transect from ~1100 m at the Chile margin site (U1542) to ~4070 m in the central South Pacific (Site U1539) and allow investigation of changes in the vertical structure of the ACC, a key issue for understanding the role of the Southern Ocean in the global carbon cycle. The sites are located at latitudes and water depths where sediments will allow the application of a wide range of siliciclastic-, carbonate-, and opal-based proxies to address our objectives of reconstructing with unprecedented stratigraphic detail surface to deep-ocean variations and their relation to atmosphere and cryosphere changes through stadial to interstadial, glacial to interglacial, and warmer than present time intervals.

Introduction

Our prior knowledge of southern high latitude paleoceanography comes mainly from conventional sediment coring and Deep Sea Drilling Project (DSDP)/Ocean Drilling Program (ODP) drilling in the Atlantic and Indian sectors of the Antarctic Circumpolar Current (ACC). A prominent example is the achievements of ODP Leg 177, which drilled sites arrayed along a north-south transect across the major ACC fronts and documented the Cenozoic history of the Atlantic sector of the Southern Ocean. These sites provided, for example, insights into changes in dust supply and marine productivity and nutrient consumption (Martínez-García et al., 2009, 2011, 2014), opposite Southern Ocean productivity between the Antarctic and Subantarctic Zone on glacial-interglacial cycles (Jaccard et al., 2013) and between the ACC and the Antarctic margin on longer timescales (Cortese et al., 2004), sea-surface temperature (SST) evo-

lution since the Pliocene (Martínez-García et al., 2010), and Antarctic Ice Sheet (AIS) dynamics during the late Quaternary (Kanfoush et al., 2002; Teitler et al., 2010, 2015; Venz and Hodell, 2002; Hayes et al., 2014).

In contrast, the Pacific sector of the ACC has been much less studied; drilling records are mostly high-resolution proximal sediment records from the Antarctic continental margin south of the ACC (e.g., Ross Sea during DSDP Leg 28 and Antarctic Peninsula during ODP Leg 178). Integrated Ocean Discovery Program Leg 318 drilled Cenozoic sediments off Wilkes Land (eastern Indian Southern Ocean) and revealed significant vulnerability of the East Antarctic Ice Sheet (EAIS) to Pliocene warming (Cook et al., 2013; Expedition 318 Scientists, 2010). A substantial advance in constraining the history and stability of the West Antarctic Ice Sheet (WAIS) and Ross Ice Shelf (RIS) came from the Antarctic Geological Drilling (ANDRILL) project (e.g., Naish et al., 2009; Florindo et al., 2008), documenting that the WAIS/RIS underwent significant changes with expanded cold and stable ice sheets between 13 and 11 Ma. The ice sheet became more dynamic in the late Miocene-early late Pliocene (10–2.5 Ma) with cyclic expansion of grounded ice sheets over the coring site after 2.5 Ma.

These on- and near-shore drilling programs have significantly advanced our understanding of AIS dynamics during the Cenozoic. However, to understand how these ice sheet changes are ultimately linked to climate, atmospheric CO₂ levels, and ocean circulation requires constraints of the ACC (i.e., the current system that connects Antarctica to the rest of the globe). In this context, our drilling in the Pacific ACC will be closely linked to the recently completed International Ocean Discovery Program (IODP) expeditions to the Ross Sea (Expedition 374), Amundsen Sea (Expedition 379), and Scotia Sea (Expedition 382). These expeditions targeted Antarctic near-shore records, which will likely be incomplete due to glacial erosion, and the potential for analyses of paleoceanographic proxies will be limited. Therefore, the Dynamics of the Pacific Antarctic Circumpolar Current (DYNAPACC; Expedition 383) pelagic sites were designed to provide a critical paleoceanographic baseline information including rates of change for improving the understanding of reconstructed Antarctic ice sheet changes and testing ice sheet models.

During Expedition 383, we recovered high-resolution Miocene to Holocene sediment records at three sites in the central Subantarctic Pacific Southern Ocean. These sites in the central South Pacific document the Miocene to Holocene ACC paleoenvironmental history at water depths from 5100 to 3600 m. We drilled two sites at the Chile margin and one site in the pelagic eastern South Pacific (ESP) to provide a depth transect (~1000–3900 m) across the major Southern Ocean water masses.

Background

The ACC is the world's largest current system and connects all three major basins of the global ocean (i.e., the Pacific, Atlantic, and Indian Oceans; Figure F1) and therefore integrates and responds to climate signals throughout the globe (e.g., Talley, 2013). By inducing pronounced upwelling and formation of new water masses, the ACC fundamentally affects the global Meridional Overturning Circulation (MOC; Marshall and Speer, 2012) and the stability of Antarctica's ice sheets.

The ACC has long been recognized as a key player in regulating atmospheric CO₂ variations and therefore global climate based on the tight coupling between Southern Hemisphere temperatures and

atmospheric CO₂ concentrations (e.g., Parrenin et al. 2013). Through its steep density surfaces, the ACC provides a direct hydrographic link between the large deep-ocean reservoir of dissolved inorganic carbon and the surface ocean along which deep-water masses may ascend to the surface (e.g., Kuhlbrodt et al., 2007; Marshall and Speer, 2012). The ACC therefore acts as a window through which ocean and atmosphere interact and as a key player in regulating atmospheric CO₂ variations and therefore global climate.

Biological utilization of nutrients in the Southern Ocean is particularly important in relation to changes in atmospheric CO₂ concentration because it regulates the preformed nutrient inventory for most of the deep ocean and therefore the global average efficiency of the biological pump (e.g., Sigman and Boyle, 2000; Sigman et al., 2010). Nutrient utilization is inefficient in the Southern Ocean today in part because phytoplankton growth is limited by the scarcity of bioavailable iron (e.g., de Baar et al., 1995). Martin (1990) proposed that dust-borne iron fertilization of Southern Ocean phytoplankton caused the ice age reduction in atmospheric CO₂. However, the role of iron in explaining variation in opal export to the Southern Ocean sediments is complex. Chase et al. (2015) found a significant correlation between annual average net primary production and modeled dust deposition but not between dust deposition and opal burial. On glacial–interglacial timescales, proxy records from the Subantarctic support a positive relationship between dust flux and opal production (e.g., Chase et al., 2003; Bratton et al., 2009; Lamy et al., 2014; Kumar et al., 1995; Anderson et al., 2014). The opposite is true south of the Antarctic Polar Front (PF) (e.g., Chase et al., 2003; Jaccard et al., 2013), indicating that factors other than dust regulate the production and export of opal in the Southern Ocean and, by association, the strength and efficiency of the biological pump. Leading candidates for these “other factors” include changes in water mass structure in the Southern Ocean locations of the fronts encircling Antarctica, which are related to the flow of the ACC, and the spatial and temporal variability of wind-driven upwelling that supplies nutrients and CO₂ to surface waters and glacial–interglacial changes in sea ice extent.

Oceanographic setting

Expedition 383 drilled sites located within the ACC current system in the central and ESP and at the southern Chile margin close to the Drake Passage. The flow of the ACC is largely driven by strong Southern Westerly Winds (SWW) and is constricted to its narrowest extent in the Drake Passage. This so-called “cold-water route” transport through the Drake Passage is one important pathway for the return of fresh and cold waters from the Pacific to the Atlantic, which strongly affects the strength of the Atlantic MOC, in concert with the “warm-water route” inflow of warm and salty Indian Ocean water masses through the Agulhas Current system (Beal et al., 2011; Gordon, 1986). The Drake Passage is ~800 km wide and located between Cape Horn and the western Antarctic Peninsula (Figure F1). Numerous hydrographic surveys across the Drake Passage since the 1970s have contributed to the understanding of seasonal and interannual variability in ACC transport through the Drake Passage and the mechanisms forcing physical and biological changes within the Southern Ocean (Meredith et al., 2011). However, even with hydrographic time series reaching back 20 y (Koenig et al., 2014) in this relatively well constrained region of the ACC and more sophisticated model simulations (Meijers, 2014), important issues such as the role of zonal winds in forcing ACC transport remain controversial. The total ACC flow transported through the Drake Passage is estimated to be between ~130 and 150 Sv (Koenig et al., 2014; Mer-

edith et al., 2011; Renault et al., 2011). The three major ACC oceanographic fronts (Orsi et al., 1995), the Subantarctic Front (SAF), the Polar Front (PF), and the Southern ACC Front (SACCF), can be identified from north to south in the Drake Passage. The modern winter sea ice margin presently does not extend significantly north into the Drake Passage. Oceanographic studies robustly show that more than half of the total Drake Passage transport occurs at and north of the SAF (i.e., in the Subantarctic Zone) followed by the Polar Frontal Zone (between the SAF and the PF). The relative contribution of Drake Passage transport south of the PF is generally <20%.

Along the southernmost Chile margin, the southward flowing Cape Horn Current (CHC), a northern branch of the ACC that continues toward the Drake Passage, provides a major fraction of the present day northern Drake Passage transport (Well et al., 2003). Satellite-tracked surface drifters reveal that after crossing the East Pacific Rise (EPR), Subantarctic surface water of the ACC is transported northeastward across the Southeast Pacific toward the Chilean coast at ~45°S, 75°W (Chaigneau and Pizarro, 2005). Here, presently only a minor part of the ACC is deflected northward into the Humboldt Current System (HCS), whereas the major fraction is deviated southward toward the Drake Passage. The CHC thus transports a significant amount of northern ACC water toward the Drake Passage in a narrow belt ~100–150 km wide off the coast (Chaigneau and Pizarro, 2005) (Figure F2).

The drill sites in the central South Pacific are all located in the Subantarctic Pacific. Sites U1539 and U1540 are located east of the EPR, whereas Site U1541 lies west of the EPR. Unfortunately, the originally proposed drill site south of the PF could not be reached during Expedition 383 because of persistent bad weather conditions. The drilled sites (U1539–U1541) lie in the pathway of the Subantarctic Pacific Antarctic Circumpolar Current, 100–170 nmi north of the modern mean position of the SAF in a zonal transition zone of the ACC. These ACC fronts form the core of the ACC and are partly constrained by the bathymetry of the EPR in this area. As a result of topographic forcing, frontal positions in the ACC close to large bathymetric features remain relatively stable at seasonal and interannual timescale, whereas over abyssal plains their location appears to be fairly unstable (Sokolov and Rintoul, 2009). The SAF at our sites at the western and eastern flank of the EPR is therefore partly constrained by topography and the location of major fracture zones (e.g., Eltanin-Tharp Fracture Zone). The oceanographic conditions in the South Pacific are influenced by the Southern Annular Mode (SAM) and the Amundsen Low but also by tropical variability and the El Niño–Southern Oscillation (ENSO) (Thompson et al., 2011; Turner et al., 2009; Yuan, 2004). Interannual and decadal SST and sea ice changes are amplified in the Pacific sector. For example, a distinct increase in sea ice north of the Ross Sea embayment and a strong sea ice decrease in the Bellingshausen–Amundsen Sea are documented by observational time series (1979–2011) (Maksym et al., 2012; Stammerjohn et al., 2012).

Deep water that upwells south of the PF feeds both Antarctic Bottom Water (AABW) close to Antarctica and Antarctic Intermediate and Mode Water formation north of the APF (e.g., Sloyan and Rintoul, 2001; Talley, 2013). The Southeast Pacific is known as the major Antarctic Intermediate Water (AAIW) formation area where fresh, cold, high-oxygen and low-nutrient waters are formed. Originally, it was thought that AAIW originates from the subduction of Antarctic surface waters below the SAF (e.g., Deacon, 1937). Later, it was suggested that the formation of AAIW is directly linked to the formation of Subantarctic Mode Water (SAMW) (McCartney, 1977), which has similar properties in the Southeast Pacific. More

recent studies basically confirm the location of the major AAIW formation area in the Southeast Pacific related to deep mixed-layer development leading to SAMW formation and finally to AAIW. The Southern Ocean's major water mass, the Circumpolar Deep Water (CDW), is divided into upper and lower branches. Lower CDW (LCDW) is identified by a salinity maximum inherited from North Atlantic Deep Water (NADW) (Reid and Lynn, 1971; Orsi et al., 1995). Upper CDW (UCDW) is typically identified as an oxygen-minimum layer derived from oxygen-depleted, nutrient-rich Indian Deep Water (IDW) and North Pacific Deep Water (NPDW) (Callahan, 1972). LCDW and UCDW are transported into the North Pacific, where they are transformed into NPDW, and return to the Southern Ocean in the East Pacific along the South American continent at 1500–3500 m water depth (Kawabe and Fujio, 2010). North of the SAF, UCDW is centered at ~1500 m depth and LCDW covers depths below 2000 m. Below LCDW, northward-spreading dense AABW formed in the Weddell Sea, Ross Sea, and along the Adélie Coast covers the abyss around Antarctica. In the South Pacific, the expansion of Ross Sea Bottom Water (RSBW), the coldest and saltiest variety of AABW, is restricted by bottom topography to areas south of the Pacific–Antarctic Ridge (Orsi et al., 1995).

Geological setting

The drill sites in the central South Pacific are all located on oceanic crust. Site U1541 is located ~160 nmi west of the crest of the EPR, north of the Eltanin–Tharp Fracture Zone. The basement age (Figure F3) (Eagles, 2006) at this location is about 10 Ma (late Miocene). Sites U1539 and U1540 are located at a similar distance to the fracture zone but ~210 nmi east of the EPR crest on slightly older crust with a middle Miocene basement age of ~15 Ma (Figure F3).

The sites at the Chile continental margin are all located on continental crust. Tectonically, the southern Chile margin is characterized by a complex geodynamic setting with oblique convergence between plates, transcurrent motion, and tectonic rotation on land (Polonia et al., 2007). The geodynamic setting of the southernmost Chile margin is driven by relative movements between three main plates: Antarctica, Scotia, and South America. This complex setting results in a strong segmentation of the southern Chile margin, which is clearly visible by bathymetric data and the available multi-channel seismic (MCS) data (Polonia et al., 2007). Two major fore-arc basins at ~53°–54.5°S and 55°–56.5°S are characterized by a thick sediment infill of as much as ~3 km (Figure F4). We drilled one site in the southern fore-arc basin (U1544). Seismic Line IT97-171 (Figure F4) documents ~3 km of remarkably undeformed sediments in this basin (Polonia et al., 2007). This southern basin most likely formed after the ridge consumption (14–10 Ma) when accretion resumed and contributed to construct the outer high that provided the barrier for the accumulation of continent-derived sediments.

North of ~53°S, subduction in the Chile trench is more orthogonal. Here, the transition between the undeformed Antarctic abyssal plain and the lower slope is very abrupt and results in a steep slope incised by canyons that run perpendicular to the trench axis. Recent sediment cover is minor at the lower slope, but a relatively small scale sediment depocenter has been localized at the upper continental slope. This sediment drift (Site U1542) contains a sediment sequence as thick as ~700 m that most likely does not extend beyond the early Pleistocene because of high sedimentation rates.

Scientific objectives

The overall goal of Expedition 383 is to improve our knowledge of Plio/Pleistocene atmosphere–ocean dynamics of the ACC in the Pacific and their implications for regional and global climate and atmospheric CO₂ based on sediment records with the highest possible resolution. The sites (Table T1; Figure F1) are located at latitudes and water depths where sediments will allow the application of a wide range of siliciclastic-, carbonate-, and opal-based proxies for reconstructing surface to deep-ocean variations and their relation to atmosphere and cryosphere changes with unprecedented stratigraphic detail. The cores recovered during Expedition 383 will allow us to test two major scientific hypotheses.

Hypothesis H1: ACC dynamics and Drake Passage throughflow conditioned the global MOC and high–low climate linkages on orbital and submillennial timescales since the Pliocene.

The Drake Passage is the major geographic constriction for the ACC and forms an important pathway for the return of upper waters to the Atlantic through the cold-water route of the MOC (Gordon, 1986). Resolving changes in the flow of circumpolar water masses through the Drake Passage is crucial for advancing our understanding of the Southern Ocean's role in affecting ocean and climate change on a global scale. Surface and intermediate water masses flowing through the Drake Passage strongly affect the Atlantic MOC together with the warm-water route inflow of Indian Ocean water masses through the Agulhas Current system (Beal et al., 2011; Knorr and Lohmann, 2003). Complementary sites below the warm-water route were drilled during IODP Expedition 361 (Hall et al., 2015). The potential role of the Drake Passage throughflow in driving changes in global MOC on glacial–interglacial and millennial timescales is not well constrained (Knorr and Lohmann, 2003; McCave et al., 2014). For the last glacial cycle, Lamy et al. (2015) reconstructed Drake Passage throughflow based on a sediment record (MD07-3128) from the Chile margin (corresponding to Proposed Site CHI-4A) and another record from the southernmost Argentinean continental slope and found a significant glacial decrease in Drake Passage throughflow (Figure F3). This finding is consistent with generally reduced current velocities of the ACC during glacials and concomitant reduction in interbasin exchange in the Southern Ocean that is most likely regulated by variations in the SWW field over the Subantarctic Zone and changes in Antarctic sea ice extent.

Downstream of the Drake Passage, a meridional section of sediment records from the Scotia Sea revealed little overall bottom current speed variations between the Last Glacial Maximum (LGM) and the Holocene (McCave et al., 2014). In contrast, a 500,000 y record from the Indian sector of the Southern Ocean suggests that the ACC was weak during warm stages and strong during glacial epochs (Mazaud et al., 2010). Likewise in the southwest Pacific, the deep western boundary current east of New Zealand intensified during glacial periods over the past 1.2 My (Hall et al., 2001), a change that has been related to a northward extension of the ACC along the New Zealand continental margin. Finally, a recent density reconstruction (based on benthic foraminifers) across the ACC south of Australia suggested a moderate increase of the ACC flow during the LGM (Lynch-Stieglitz et al., 2016). Taken together, there is substantial disagreement about the ACC flow speed and its po-

tential impact on changes in Drake Passage throughflow in response to glacial and interglacial cycles, which prevents us from a more complete understanding of MOC dynamics.

Superimposed on the long-term glacial reduction of Drake Passage throughflow, prominent high-amplitude, millennial-scale variability was observed, implying a strong sensitivity of the northern ACC in the Southeast Pacific to suborbital climate variations (Lamy et al., 2015). Enhanced Drake Passage throughflow during Antarctic warm periods is consistent with the previously suggested impact of the bipolar seesaw mechanism on the Southern Ocean (e.g., Anderson et al., 2009; Lamy et al., 2007), which leads to, for example, surface water warming, enhanced upwelling, and a stronger ACC caused by southward-shifted SWW. The glacial reduction of the cold-water route occurred in concert with the well-documented decrease of the warm-water route (Aguilhas leakage; Beal et al., 2011). The data currently available thus indicate that both of these oceanographic corridors are critical for the South Atlantic contribution to glacial MOC strength changes.

Our sites at the Chile margin (U1542 and U1544) and in the ESP (U1543) (Figure F2) provide a unique opportunity to extend the existing 60 ky record back into the early Pleistocene and Pliocene and address important questions such as the role of long-term, orbital-scale, and millennial-scale changes in ACC strength and Drake Passage throughflow in warmer-than-present interglacials (e.g., Marine Isotope Stages [MISs] 5e, 11, and 31) or the Pliocene warm period and the link to Antarctic climate fluctuations over multiple glacial–interglacial cycles.

In the Pacific sector of the Southern Ocean, atmosphere–ocean changes between the high, middle, and low latitudes are strongly linked. Pronounced past changes in surface–ocean properties at both millennial and orbital timescales, particularly SSTs, indicate an exceptional climate sensitivity to external and internal forcings on these timescales. Glacial–interglacial amplitudes in Subantarctic SSTs are generally $>4^{\circ}\text{C}$ in the Atlantic (Gersonde et al., 2005; Martínez-García et al., 2010), Indian (e.g., De Deckker et al., 2012; Lorrey et al., 2012), and Pacific Southern Ocean (Caniupán et al., 2011; Ho et al., 2012; Pahnke et al., 2003). Millennial-scale fluctuations with amplitudes of $\sim 2^{\circ}$ – 3°C are observed at continental margin sites in the Subantarctic Pacific during the last glacial (Barrows et al., 2007; Caniupán et al., 2011; Lamy et al., 2004; Pahnke et al., 2003).

High-amplitude glacial–interglacial SST changes in the Subantarctic Pacific Southern Ocean (Ho et al., 2012) are closely coupled to those observed in the subtropical ESP, implying large shifts of the northern ACC system and export of Subantarctic surface water masses into the HCS. This variability is probably directly related to variations in the strength and position of the ACC fronts, the SWW, and Drake Passage throughflow (Lamy et al., 2015) (Figure F5).

Along the Chile margin, the reconstructed glacial cooling implies a stronger northward extension of the Antarctic cold-water influence in the ESP during glacials (Ho et al., 2012) compared to the central Pacific (Figure F3). The significant glacial decrease in Drake Passage transport accompanied by a reduction of the ACC transport strongly reduced the interbasin water mass exchange in the Southern Ocean and enhanced the high–low latitude exchange in the Pacific Eastern Boundary Current system (Figure F3). The reacceleration of subantarctic Drake Passage transport matches the major deglacial warming in the Subantarctic Southeast Pacific and South Atlantic (Barker and Diz, 2014), suggesting a close coupling of Drake Passage throughflow to the initiation of Southern Hemisphere warming.

Similar to the glacial–interglacial pattern, the inferred millennial-scale changes in Drake Passage throughflow (Lamy et al., 2015) are paralleled by SST changes in proxy records from the Southeast Pacific (ODP Site 1233) (Kaiser and Lamy, 2010) that extend north to the cold tongue in the eastern tropical Pacific (Lea et al., 2006). These data imply that substantial changes in the oceanographic dynamics of the South Pacific Gyre are related to the northward deflection of ACC waters at millennial timescales. Paleoceanographic reconstructions based on our sites at the Chile margin (U1542 and U1544) and in the ESP (U1543) in combination with ODP Leg 202 sites (Figure F2) (e.g., Dekens et al., 2007; Lamy et al., 2004; Pena et al., 2008; Rincon-Martinez et al., 2009) and the JR100 expedition (http://iodp.tamu.edu/scienceops/expeditions/JR100_379T.html) sites will allow us to estimate changes in the coupling between the ACC and the tropical Pacific over various timescales and at warmer-than-present climate states. High–low latitude linkages (both modern and past) in the oceanographically little explored ESP are important for global climate. The suggested enhanced glacial northward export of cold-water masses to the tropical Pacific reinforces earlier modeling studies showing the importance of southerly derived surface and intermediate water masses in the tropical Pacific for global cooling during the LGM (Liu et al., 2002).

Our drill sites will allow us to better constrain the latitudinal shifts of the Southern Ocean fronts over time to understand the resulting changes in ACC transport, intrabasin mixing, and AAIW production over a wider range of climate states. For example, reconstructions of past changes in the latitudinal position of the SAF suggest as much as 9° latitude northward shifts during glacials, whereas other areas suggest no shift at all (Kohfeld et al., 2013). Smaller shifts probably occurred where the front is bathymetrically confined, like in the vicinity of the EPR (central South Pacific sites). Large shifts likely occurred in the eastern Subantarctic Pacific with the SAF potentially located north of the Drake Passage during peak glacials (Gersonde et al., 2003; Ho et al., 2012; Lamy et al., 2015). As stated above, these changes may substantially affect the Drake Passage throughflow and low–high latitude exchange of heat and of nutrients with major implications for the global climate development. Further south, Benz et al. (2016) reconstructed a glacial winter sea ice (WSI) expansion by $\sim 2^{\circ}$ – 3° in latitude in the western part of the Pacific sector and an even stronger shift ($\sim 5^{\circ}$ in latitude) of both the SACCF and the WSI edge in the central part (Figures F6, F7). This west–east differentiation in the winter SST pattern and sea ice extent can be related to differences in topographic and atmospheric forcing (Benz et al., 2016).

Despite the strong climate sensitivity of the Subantarctic Southern Ocean, only one SST record is available (ODP Site 1090 in the Subantarctic Atlantic) that reaches back to the Pliocene (Figure F8). Here, the major cooling phase after ~ 1.6 Ma coincides with the emergence of the eastern tropical Pacific cold tongue and an intensification of the meridional temperature gradients (Martínez-García et al., 2010) (Figure F4). The ANDRILL records suggest that high-latitude climate cooling between 3.0 and 2.5 Ma drove both the WAIS and EAIS toward their present expanded cold polar state (Naish et al., 2009). The intensification of Antarctic cooling resulted in strengthened SWW and northward migration of Southern Ocean fronts, which would restrict water exchange between ocean basins (reduced MOC) beginning after 3.3 Ma (McKay et al., 2012). Warm early Pleistocene and Pliocene SSTs are consistent with the poleward expanded warm pool (Brierley et al., 2009)

Our sites from the central South Pacific (U1539–U1541) will allow a more comprehensive assessment of the temperature evolution of the Pacific Southern Ocean and its relation to atmospheric variations of the SWW, sea ice, and AIS variability during cold and warmer-than-present (e.g., MISs 5, 11, and 31 and the Pliocene warming) time periods. Information on warm periods is critical to assess the Southern Ocean's role under future warming, especially for AIS melting and associated sea level rise because ice sheet modeling together with modern observations show that oceanic forcing is the most important forcing mechanism (Pollard and DeConto 2009; Paolo et al., 2015).

Moreover, temperature records from our high-resolution sites at the Chile margin (U1542 and U1544) and central South Pacific (U1539 and U1540) will allow us to compare millennial-scale variations between the Pacific Southern Ocean and Antarctica for a time interval that is well beyond the reach of ice cores. Even on a global scale, such millennial-scale temperature records back into the Pliocene are rare (e.g., IODP Site U1313: Naafs et al., 2013; ODP Site 980: McManus et al. 1999).

Hypothesis H2: Variations in the Pacific ACC determine the physical and biological characteristics of the oceanic carbon pump and atmospheric CO₂.

Atmosphere-ocean-cryosphere interactions and teleconnections between high and low latitudes play an important role in understanding processes and feedbacks of past and future climate change, and the Subantarctic Southern Ocean provides the major link between Antarctica and the low latitudes. In the Southern Ocean, these interactions are believed to control sea ice cover, AIS dynamics, upper ocean stratification, biological nutrient utilization, and exposure rates of deep water. They have been considered to play a key role in explaining the variability in atmospheric CO₂ concentrations. At low latitudes, oceanic circulation changes have profound impacts on large-scale atmospheric circulation patterns, particularly in regard to trade and westerly winds. The SWW strongly impact the upwelling of carbon-rich deep-water masses in the Southern Ocean affecting atmospheric CO₂ (e.g., Anderson et al., 2009) and also control the return flow of intermediate waters to the tropics (Figure F9). Moreover, the SWW exert an important control on the upwelling of relatively warm deep water onto the West Antarctic continental shelf in the Pacific sector where it causes sub-ice shelf melting and mass loss of the WAIS (e.g., Steig et al., 2012). Thus, the SWW are thought to play a key role in past and future climate change, and yet their strength and position in colder and warmer climates relative to today remain a wide-open question (Kohfeld et al., 2013).

In the modern Southern Ocean, deep waters with high dissolved inorganic carbon (DIC) and major nutrient concentrations are carried to the surface by wind-driven upwelling and density-driven overturning. However, phytoplankton growth is limited by the scarcity of iron, and these nutrients return to the subsurface before they are completely consumed. This incomplete utilization of nutrients represents a substantial leak in the modern global biological pump because it allows the escape of deeply sequestered carbon back to the atmosphere, keeping atmospheric CO₂ levels high during interglacial stages. An increase in the efficiency of the global biological pump can be accomplished either by decreasing the physical cycling of deep water through the surface of the Southern Ocean or by increasing the degree to which surface nutrients are consumed by marine organisms (e.g., as a result of increasing iron availability relative to other nutrients) (Sigman and Boyle, 2000; Sigman et al., 2010).

Paleoceanographic records (mainly from the Atlantic sector of the Southern Ocean) indicate that ice age productivity was lower (Anderson et al., 2009; Kohfeld et al., 2005; Jaccard et al., 2013) in the Antarctic zone (i.e., south of the PF) and nutrient consumption was higher (Robinson and Sigman, 2008; Studer et al., 2015). These data indicate that the Antarctic zone was more strongly stratified during ice ages, reducing the Antarctic leak in the biological pump and therefore contributing to the lowering of atmospheric CO₂ during glacial times.

In the Subantarctic Atlantic, higher dust flux, increased productivity and nutrient consumption, and lower atmospheric CO₂ covary during peak glacial times and millennial cold events, which has been associated with iron fertilization (e.g., Frank et al., 2000; Kumar et al., 1995; Martínez-García et al., 2009, 2011, 2014). Martínez-García et al. (2009, 2014) suggested that around 40–50 ppmv of the glacial atmospheric CO₂ decline can be attributed to iron fertilization (Figure F10), which is in good agreement with model estimates (Hain et al., 2010; Chalk et al., 2017; Brovkin et al., 2007). At ODP Site 1090, the first significant rise in ice age dust and iron deposition over the past 4 My occurred at ~2.7 Ma, and the most dramatic rise in dust deposition occurred across the mid-Pleistocene transition (Martínez-García et al., 2011).

For the past million years, higher dust deposition during glacial periods compared to interglacials has also been documented for the Pacific Southern Ocean, suggesting an eastward extension of the influence of Australian and/or New Zealand dust sources (Lamy et al., 2014). In this area, dust fluxes follow the same glacial–interglacial fluctuations, but absolute glacial fluxes in the Subantarctic Pacific are about half of those in the Atlantic Southern Ocean (Figure F11). As in the Atlantic sector, export production in the Subantarctic Zone of the Pacific sector over the past ~450 ky is higher during glacial times, but in contrast to the Subantarctic Atlantic (Martínez-García et al., 2014), the relationship between productivity and nutrient consumption has not been evaluated in detail yet. Our preliminary data suggest that nutrient consumption increased in response to iron fertilization (A. Martínez-García, unpubl. data).

The drill cores from the Pacific sector of the Southern Ocean will allow exploration of the glacial–interglacial changes in biological productivity both north and south of the Antarctic PF in more detail. Core records, including those from ODP Leg 177, show that the glacial-time biological productivity in the Subantarctic Zone of the Atlantic sector of the Southern Ocean increased during the glacial periods of the late Quaternary and decreased in the Antarctic Zone of this sector (Diekmann, 2007; Jaccard et al., 2013). The glacial productivity decrease in the Antarctic Zone is under debate and may have been caused either by stratification of the water column (Sigman et al., 2004; Jaccard et al., 2005) or expansion of Antarctic sea ice coverage (Hillenbrand and Cortese, 2006). The new drill core records proposed here will allow us to test whether this pattern is also observed in the Pacific sector of the Southern Ocean, as indicated by Bradtmiller et al. (2009), and will help to identify the cause for the potential glacial productivity decrease south of the PF.

Our drill sites at the central South Pacific (U1539–U1541) will provide robust data to test to which extent processes that are active in the Atlantic sector can be translated to the Pacific sector, thus allowing the construction of a more global picture of the Southern Ocean's role in nutrient distribution and biogenic export production and their impact on CO₂ variations. It is necessary to solve these issues before the impact on the ocean carbon cycle under different climate boundary conditions can be modeled more realistically.

Moreover, the ESP site (U1543) and Chile margin sites (U1542 and U1544) will provide a long-term record of variability of the Patagonian Ice Sheet (PIS). Both terrestrial records from Patagonia and records of ice-rafted debris (IRD) in marine sediment cores from the Chile margin (MD07-3128 at Proposed Site CHI-4A) indicate substantial millennial-scale variability of the PIS paralleling dust input changes recorded in Antarctic ice cores. This covariance implies a causal link between ice sheet advances and dust availability (Caniupán et al., 2011; Sugden et al., 2009). Thus, the Chile margin sites will provide important baseline information for dust variability in the Atlantic sector of the ACC.

Our two hypotheses contain a large number of challenging research targets critical for paleoceanography and paleoclimatology on a global scale. Our sites in the central South Pacific represent the first drill cores in the Subantarctic Pacific sector of the Southern Ocean. Through comparison to the ESP site, we include a particularly sensitive ocean area where glacial/interglacial changes are strongly amplified. Finally, the Chilean Margin Sites provide information on changes at shallower water depths together with changes in the PIS and potential links to dust input in the Atlantic sector of the Southern Ocean.

Site summaries

Site U1539

Background and objectives

Site U1539 (Proposed Site CSP-2B) is located in the central South Pacific at 56°09.0655'S, 115°08.038'W ~1600 nmi west of the Magellan Strait at 4070 m water depth (Figure F12). The site sits at the eastern flank of the southernmost EPR, ~220 nmi from the modern seafloor spreading axis, and is underlain by oceanic crust formed at ~10–12 Ma (Eagles, 2006). Assuming a constant seafloor half-spreading rate of ~4.5 cm (Pitman and Heirtzler, 1966), the plate tectonic backtrack path of Site U1539 moves the site westward. This translates to an early Pliocene position ~100 nmi closer to the crest of the EPR at a water depth shallower by several hundred meters. At a smaller scale, the site is located in a northeast–southwest oriented, ~5 nmi wide trough that parallels the orientation of the EPR. The adjacent ridges rise to ~3000 m water depth northwest of the site and to ~3500 m water depth to the southeast (Figure F12).

Site U1539 is located on MCS Line AWI-201000013 (Figure F13A) close to the intersection with Line AWI-201000011 (Gersonde, 2011). The seismic cross-lines indicate ~650 m thick sediments above oceanic basement. Sediments are mostly well stratified with flat-lying reflectors. Low to moderately reflecting layers become stronger deeper than ~100–120 m sediment depth. Sediment echo sound (Parasound) profiles (Gersonde, 2011) reveal excellent penetration (>150 m) with distinct layering (Figure F13B), suggesting a succession of fine-grained soft sediments with varying lithologic composition.

Site U1539 lies in the pathway of the SAP ACC, ~100 nmi north of the modern average SAF, in a zonal transition zone of the ACC. West of the site, the ACC and associated fronts are strongly steered by the topography of seafloor spreading systems (Udintsev and Eltanin-Tharp Fracture Zone systems), whereas to the east the vast Amundsen Sea basin does not influence the ACC strongly.

SSTs seasonally vary between ~2°C (July–September) and ~6.5°C (January–March). The area is located west of the main Antarctic Intermediate and Mode Water formation regions in the Southeast Pacific. The 4070 m water depth places Site U1539 within

LCDW, close to the boundary to AABW (Figure F14). This hydrographic setting makes it ideal to evaluate past changes in frontal position, associated export production, ACC current speed and position, and eolian dust and IRD input during the Pleistocene.

The main objectives at Site U1539 were to

- Recover a moderate- to high-resolution Subantarctic Pliocene–Quaternary sediment record close to the SAF;
- Investigate the sequencing of siliceous and calcareous oozes, allowing for a wide range of paleoceanographic reconstructions;
- Reconstruct high-amplitude Subantarctic SSTs and sea ice variations;
- Provide a record of lowermost CDW and glacial AABW;
- Reconstruct productivity (opal versus carbonate), nutrient distribution, and dust-productivity coupling; and
- Recover a potential far-field record of WAIS variability.

Operations

The first core of Expedition 383 was taken in Hole U1539A at 0445 h on 1 June 2019 from 4071.1 meters below sea level (mbsl). We used the advanced piston corer (APC) system to penetrate from the seafloor to 107.6 meters core depth below seafloor, Method A (CSF-A) (Cores 383-U1539A-1H through 12H) and recover 104.75 m of core (97%). The hole was terminated after Core 12H because of rough weather and sea conditions. Shattered liners were recorded on Cores 5H and 11H. Shear pins were sheared prematurely on Cores 1H, 11H, and 12H.

After 27 h of waiting on weather, Hole U1539B was spudded at 0540 h on 3 June. Cores 383-U1539A-1H through 3H recovered 21.9 m (78% recovery), but Cores 2H and 3H were misfires, so we decided to terminate the hole and wait for seas to further improve before continuing operations.

By 1245 h, heave had fallen to 3.0 m and we resumed coring. Hole U1539C was started with Core 383-U1539C-1H returning a full 9.7 m core barrel. This allowed us to determine a seafloor depth of 4070.2 mbsl. Full-length APC coring continued to refusal at 240.1 m CSF-A (Core 26H). Half-length APC (HLAPC) coring was then used to deepen the hole, and coring terminated after Core 32F at 268.1 m CSF-A. A total of 32 cores were taken in this hole using the APC and HLAPC coring tools. Formation temperature measurements were taken with Cores 4H, 13H, and 16H. Partial strokes were recorded on Cores 24H–27F with high overpull on Cores 17H–26H. A misfire was recorded on Core 31F.

Hole U1539D was spot-cored to fill in coring gaps from the previous holes. The hole was spudded at 1405 h on 5 June, drilled ahead without recovery to 47.5 m CSF-A, where coring began, and continued to 198 m CSF-A with seven drilled intervals totaling 68.7 m. A total of 14 APC cores were taken over the 129.3 m cored interval with a recovery of 128.15 m (99%). The hole reached maximum depth at 1500 h on 6 June, and the drill string was recovered, ending the hole and Site U1539. Misfires were recorded on Cores 383-U1539D-2H, 3H, 6H, 8H, and 9H, and a partial stroke was recorded on Core 18H. No temperature or orientation measurements were taken in this hole. After recovering the drill string, the rig floor was secured for transit at 0130 h. The thrusters were raised, and the sea voyage to Site U1540 began at 0224 h on 7 June.

Principal results

At Site U1539, a 268.1 m thick continuous sequence of Holocene to early Pleistocene sediments was recovered with a bottom-hole age of ~1.3 Ma and a mean sedimentation rate of ~20.6 cm/ky. Hole U1539C recovered the deepest cored sediments at 268.1 m CSF-A.

The recovered 247.50 m of core spans one lithologic unit with two subunits consisting of interbedded calcareous and siliceous biogenic ooze (Figure F15). Site U1539 sediments can broadly be divided into two categories: diatom ooze with varying amounts of calcareous and biosiliceous components and biogenic carbonate ooze with varying amounts of biosiliceous and other calcareous components. Siliceous microfossils show relatively good preservation and are abundant throughout the record. Preservation of calcareous microfossils varies from moderate to good.

Five primary lithofacies were identified, and they mostly differ in their proportions of calcareous and biosiliceous components. Secondary defining attributes include degree of diagenetic alteration, bioturbation, and in the case of Lithofacies 5, microfossil size and sedimentary structure. Lithofacies 1 comprises distinct centimeter-scale diatom mats in a biogenic ooze matrix. This facies can be distinguished from Lithofacies 2, which is characterized by sequences of more pronounced and stacked diatom mats with a near absence of diatom ooze matrix. Lithofacies 3 is the dominant lithofacies at Site U1539. It consists of light greenish gray to gray carbonate-bearing or carbonate-rich diatomaceous ooze that exhibits moderate to heavy bioturbation. Lithofacies 4 consists of white to light gray diatom-bearing to diatom-rich nannofossil or calcareous oozes that occur as thin beds generally <3 m thick with moderate to heavy bioturbation. Lithofacies 5 corresponds to foraminiferal oozes and sands that occur sporadically as centimeter- to decimeter-scale layers characterized by a sharp basal contact and distinct cross-bedding observable in X-ray images.

We divided the Site U1539 stratigraphic sequence into Subunits IA and IB based on the relative abundance of the five defined lithofacies. With the exception of Lithofacies 5 in Subunit IB, all lithofacies are present in both subunits, but the thickness, frequency, and overall occurrence differs between the two. Subunit IA is characterized by the frequent occurrence of ~1.5 m thick beds of calcareous and nannofossil ooze (Lithofacies 4) and limited occurrence of diatom mats (Lithofacies 2). In contrast, Subunit IB has only a few occurrences of Lithofacies 4 and is largely dominated by beds of thick and continuous diatom mats (Figure F16). Carbonate-bearing and carbonate-rich diatomaceous ooze (Lithofacies 3) is the dominant lithofacies in both subunits and does not provide a useful means to divide the stratigraphy because of its widespread occurrence. The prominent white to light gray calcareous beds associated with Lithofacies 4 drive the high-amplitude variations in RGB blue and L^* values. In general, we observed high-amplitude, low-frequency variations above the 125 m core composite depth below seafloor, Method A (CCSF-A), subunit boundary and more frequent low-amplitude variations below it (Figure F16).

Biostratigraphic age assignments are consistent with each other and are mainly derived from diatom, radiolarian, and calcareous nannofossil datums. Diatom, radiolarian, and nannofossil biostratigraphy detected no major hiatuses, indicating that a continuous sediment sequence from the early Pleistocene through the Holocene was recovered at this site.

We could not observe unambiguous evidence for geomagnetic reversals in any of the holes at Site U1539. This is probably due to low natural remanent magnetization (NRM) intensities that could be related to the high diatom content of the sediments. Therefore, the paleomagnetic record cannot contribute to the age model for Site U1539.

We analyzed samples for headspace gas, interstitial water chemistry, and bulk sediment chemistry at a resolution of three per core in Hole U1539A and three per core from 78.9 to 120 m CSF-A, two

per core from 120 to 150 m CSF-A, and one per core from 150 m CSF-A to the bottom of Hole U1539C at 266.41 m CSF-A. Headspace samples were not collected from HLAPC cores. Methane concentration is low at this site overall. It gradually increases downhole, averaging 12.3 ppmv and never exceeding 18.4 ppmv. Ethane and propane remain below detection limit throughout the entire hole. In Hole U1539A, alkalinity increases in the uppermost 10.6 m CSF-A to ~5 mM, likely caused by anaerobic organic matter remineralization. pH values also increase from 7.7 at the surface to ~8 at 2.94 m CSF-A and remain constant with depth. Interstitial water magnesium and lithium concentrations appear conservative throughout the sediment column. Dissolved manganese concentrations increase sharply from below detection limit in the mudline sample to a maximum of 60 μ M at 7.57 m CSF-A, indicating anaerobic, reductive dissolution of manganese oxides within the uppermost 3 m of sediment.

The composite carbonate record at Site U1539 (Holes U1539A and U1539C) shows downhole variability ranging from 2.9 to 92.8 wt% and a strong correlation with RGB blue and reflectance L^* measurements (Figure F17) and gamma ray attenuation (GRA) bulk density. Total organic carbon (TOC) concentration has a mean value of 0.40 wt%, and values range between a minimum of 0.14 wt% and a maximum of 0.76 wt% (Hole U1539C). The mean value of TOC/total nitrogen (TN) is often >10 from 0.8 m CSF-A in Hole U1539A and to 115.2 m CSF-A in Hole U1539C, suggesting that the location of Site U1539 received a fraction of terrestrial organic matter input across these depths.

Physical properties measurements at Site U1539 included non-destructive whole-round measurements of GRA bulk density, magnetic susceptibility (MS), Whole-Round Multisensor Logger (WRMSL) P -wave velocity, and natural gamma radiation (NGR) on core sections from Holes U1539A–U1539D. Additional physical properties collected include thermal conductivity (TCon) on whole-round cores from all holes. Downhole changes in physical property characteristics overall are in good agreement with the lithofacies defined based on sedimentologic characteristics (Figure F18). In general, diatom ooze lithofacies (1 and 2) correspond to high MS, low bulk density, and high porosity (Figure F18). The lowest densities and highest porosities occur in diatom mats within diatom ooze, which is the recurring dominant Lithofacies 2. In contrast, intervals that are predominantly calcareous nannofossil ooze (Lithofacies 4 and 5) correspond with decreased NGR, high bulk density, low P -wave velocity, low to moderate MS, and low to moderate porosity. Over most of the record, GRA bulk density shows an anticorrelation with the NGR-derived potassium concentrations and MS. The NGR record also exhibits 10–15 m scale cyclicity, particularly in the interval from ~10 to 135 CSF-A. NGR data also imply that total counts and the derived potassium concentrations can be used as a semiquantitative proxy for the abundance of terrigenous material delivered by dust or sea ice and iceberg transport.

Correlations between holes at Site U1539 were accomplished using the Correlator software (version 3.0). We constructed a splice for the upper portion of the site from 0 to 219.80 m CCSF-A using Holes U1539A–U1539D. However, the splice contains several gaps, as detailed below. Deeper than 219.80 m CCSF-A, only a single hole (U1539C) was drilled; thus, Cores 383-U1539C-22H through 32F were appended to the splice with gaps set between each core to the bottom of the hole. The composite depth scale is anchored to the mudline of Core 383-U1539A-1H, which is assigned as 0 m CCSF-A.

We constructed a preliminary age model based on biostratigraphic results, mainly including diatom and nannofossil age markers. These data suggest that the sedimentary sequence recovered at Site U1539 covers the past ~1.3 My (Figure F19). For a pelagic setting, sedimentation rates are extraordinarily high, reaching on average 17 cm/ky back to 0.9 Ma and even higher values of ~30 cm/ky from ~0.9 to 1.3 Ma. This biostratigraphic age model is generally consistent with preliminary stratigraphic tuning performed on-board based on physical property data such as color measurements (RGB blue) and GRA bulk density.

The combination of nearly continuous recovery, very high sedimentation rates driven by high diatom productivity and potential sediment focusing, clear patterns in physical properties and sediment color, and a rich array of well-preserved diatoms combined with calcareous microfossils will provide unprecedented opportunities for improving our understanding of the dynamics of the ACC and its link to global carbon cycle changes at orbital and suborbital timescales.

Site U1540

Background and objectives

Site U1540 (Proposed Site CSP-7A) is located in the central South Pacific at 55°08.467'S, 114°50.515'W ~1600 nmi west of the Magellan Strait at 3580 m water depth (Figure F20). The site sits at the eastern flank of the southernmost EPR within the Eltanin-Tharp Fracture Zone, ~130 nmi from the modern seafloor spreading axis, and is underlain by oceanic crust formed at the EPR about 6–8 Ma ago (Eagles, 2006). Assuming overall constant seafloor half-spreading rates of ~4.5 cm/ky (Pitman and Heirtzler, 1966), the plate tectonic backtrack path of Site U1540 moves the site westward to an early Pliocene position ~100 nmi closer to the crest of the EPR at a water depth shallower by several hundred meters. At a smaller scale, the site is located at the northeast end of a ridge that parallels the orientation of the EPR.

Site U1540 is crossed by a sediment echo sound (Parasound) profile that is oriented northwest–southeast. The Parasound profile documents excellent penetration (>150 m) with distinct layering (Figure F21), suggesting a succession of fine-grained soft sediments with varying lithologic composition. The site is also located ~2 nmi northwest of site survey Core PS75/56 (Gersonde, 2011) (Figure F5).

Site U1540 lies in the pathway of the SAP ACC, ~170 nmi north of the modern mean position of the SAF in a zonal transition zone of the ACC. West of the site, the ACC and the associated fronts are strongly steered by the topography of seafloor spreading systems (Udintsev and Eltanin-Tharp Fracture Zone systems), whereas to the east the vast Amundsen Sea basin does not influence the position of the ACC strongly.

SSTs seasonally vary between ~2.5°C (July–September) and ~7°C (January–March). The area is located west of the main Antarctic Intermediate and Mode Water formation regions in the Southeast Pacific. The 3580 m water depth places Site U1540 within LCDW (Figure F14). This hydrographic setting makes it ideal to evaluate past changes in frontal position, associated export production, ACC current speed and position, and eolian dust and IRD input during the Pleistocene and Pliocene.

The main objectives at Site U1540 were to

- Recover a moderate- to high-resolution Subantarctic Pliocene–Quaternary sediment record close to the SAF;

- Investigate the sequencing of siliceous and calcareous oozes, allowing for a wide range of paleoceanographic reconstructions;
- Reconstruct high-amplitude Subantarctic SSTs and sea ice variations;
- Provide a record of LCDW and glacial AABW;
- Reconstruct export production (opal versus carbonate), nutrient distribution, and dust-productivity coupling; and
- Recover a potential far-field record of WAIS variability.

Operations

The sea voyage to Site U1540 began at 0224 h on 7 June 2019. It was a short 61 nmi transit that was completed in only 6.1 h, and the ship arrived on site at 0830 h. The thrusters were lowered, and the vessel was in full dynamic positioning mode over the site coordinates by 0929 h. The drill crew made up an APC/extended core barrel (XCB) bottom-hole assembly (BHA) and lowered the drill string to the seafloor, placing the bit at 2171 meters below the rig floor (mbrf) by 1645 h. The vessel began experiencing heave above 7 m with a roll of ~5° at that time, and it was decided to wait for the seas to calm before continuing operations.

By 0415 h on 8 June, the seas had calmed sufficiently to continue operations. With an estimated depth of 3600.6 mbrf (calculated by the precision depth recorder [PDR]), the bit was lowered to 3595.0 mbrf to spud Hole U1540A. Core 383-U1540A-1H recovered 9.0 m of sediment and determined a seafloor depth of 3584.6 mbsl. Hole U1540A was deepened to 150 m CSF-A, and Cores 1H–16H recovered 155.1 m (103%) by 0815 h on 9 June. The top drive was set back, and the bit was pulled to 3561 mbrf, clearing the seafloor at 0900 h on 9 June and ending Hole U1540A. Formation temperature measurements were taken with the advanced piston corer temperature tool (APCT-3) on Cores 4H, 7H, 10H, 13H, and 16H, and orientation measurements were taken on all cores. Misfires were recorded on Cores 1H, 7H, and 16H. A misfire on Core 7H caused an unreliable temperature reading.

The vessel was moved 20 m east, and the bit was lowered to 3575 mbrf to spud Hole U1540B. Our first attempt to take a mudline core did not retrieve any sediment. The bit was then lowered 9.5 m to 384.5 mbrf, and Hole U1540B was started at 1220 h on 9 June. Based on the recovery in Core 383-U1540B-1H, the seafloor was calculated at 3580.0 mbsl. APC coring continued to 150 m CSF-A with three drilled intervals (totaling 10 m) to help obtain good overlap with cores from Hole U1540A. A total of 16 cores were taken over a 140.8 m interval with 101% recovery. Misfires were recorded on Cores 1H, 6H, 8H, and 11H. The bit was raised, clearing the seafloor at 0958 h on 10 June and ending Hole U1540B.

After coring Hole U1540B, permission was requested from the IODP Environmental Protection and Safety Panel to extend the allowed penetration depth at Site U1540 from 150 to 275 m CSF-A. This request was based on the successful recovery of a complete and continuous Pleistocene sedimentary record in Holes U1540A and U1540B. The increase in penetration depth would allow us to extend this record into the Pliocene, which was expected directly below the already cored maximum depth of 150 m CSF-A. An additional reason for the request was the prevailing sea conditions, which predicted two additional days of reasonably calm seas, thus allowing for further drilling, whereas significantly worst sea conditions were forecast at our two other primary sites in the South Pacific.

Hole U1540C was spudded 20 m south of Hole U1540B at 1110 h on 10 June with the bit positioned at 3587.5 mbrf. Core 383-

U1540C-1H recovered 6.8 m, but the top of the core lacked the characteristic light brown oxidized layer that would have indicated the recovery of a good sediment–water interface. Therefore, we decided to end Hole U1540C and attempt another mudline core.

We repositioned the vessel 20 m west of Hole U1540C and started Hole U1540D with the bit at 3587.5 mbrf. Core 383-U1540D-1H recovered 8.9 m (100%) with a noticeable light brown sediment layer at the top of the core. The amount of core recovered determined a water depth of 3577.5 mbsl. APC coring continued to 79.2 m CSF-A with three interspersed, short drilled intervals to ensure adequate core overlap for stratigraphic correlation between Holes U1540A, U1540B, and U1540C.

Permission to extend the penetration depth at Site U1540 to 275 m CSF-A was received while retrieving Core 383-U1540D-11H from 79.2 m CSF-A. At that point, we drilled ahead without recovery to 131.0 m CSF-A before resuming coring operations. The APC coring system was redeployed, and the hole was deepened until we reached APC refusal at 210.5 m CSF-A. Cores 21H and 22H (207.0–210.5 m CSF-A) recovered just 3.39 and 0.5 m of sediment, respectively, with numerous rock fragments. We chose to deploy the XCB system for one more core in an attempt to recover more of the presumed basement material, but after coring for 45 min with only 0.5 m of advancement, the core barrel was pulled back on board with no recovery. We raised the bit, which cleared the seafloor at 1710 h on 11 June, ending Hole U1540D. A total of 19 cores were taken over a 151.8 m interval with 151 m recovered (99%). Misfires were recorded on Cores 4H and 6H, and partial strokes were recorded on Cores 22H and 23H.

The vessel was offset another 20 m west of Hole U1540D, and the bit was spaced out for spudding another hole using the seafloor depth calculated from Hole U1540D. Hole U1540E was spudded at 1840 h on 11 June and drilled ahead without recovery to 135.0 m CSF-A. At that point, we deployed the APC system and cored continuously until APC refusal at 213.3 m CSF-A (Cores 383-U1540E-2H through 10H). The pipe was recovered on board, and the bit cleared the rotary table at 1900 h on 12 June. The rig was secured and readied for transit by 1905 h. A total of eight cores were taken in Hole U1540E with the APC system over a 76.0 m interval with 78.9 m recovered (104%). Core 10H experienced a partial stroke. Two drilled intervals advanced 137.0 m.

With the seas too high to raise the thrusters, the vessel waited on weather until the seas calmed down at 1545 h on 13 June. The thrusters were then raised, and the vessel began the transit to Site U1541.

Principal results

A ~213 m thick continuous sequence of Holocene to early Pliocene sediments was recovered at Site U1540 (Figure F22). A bottom-hole age of ~4.8 Ma and a mean sedimentation rate of ~4.5 cm/ky were determined. The deepest cored sediments were recovered in Hole U1540E. A spliced sedimentary sequence ~227 m in length was constructed from Holes U1540A–U1540E. Five primary lithofacies were identified at this site (Figure F23). The numbering of facies is based on all facies documented for Expedition 383 so far. The sequence is dominated by light greenish gray to gray carbonate-bearing to carbonate-rich diatom oozes (Lithofacies 2). These are frequently interbedded with light gray or light greenish gray diatom-bearing to diatom-rich nannofossil or calcareous oozes (Lithofacies 3). Less abundant are white to very light gray nannofossil oozes of heavily bioturbated, massive appearance (Lithofacies 4) and greenish gray to gray clay-bearing to clay-rich biogenic oozes

(Lithofacies 6 and 7). Based on the distribution and co-occurrence of the defined lithofacies, we divided the Site U1540 sedimentary sequence into two lithostratigraphic units and the upper Unit I into two subunits, IA and IB. Subunit IA extends downhole to ~58 m CSF-A (~62 m CCSF-A) and is dominated by Lithofacies 2 and thin intercalated beds of Lithofacies 4. It is therefore broadly equivalent to Subunit IA at Site U1539. Subunit IB spans from ~58 to ~156 m CSF-A (~62 to 178 m CCSF-A) and shows an increased occurrence of Lithofacies 3 and a decreased contribution of Lithofacies 2. Lithofacies 1, 4, and 6 are virtually missing in this subunit. Unit II extends from 178 m CCSF-A to the base of the record at 227.1 m CCSF-A (165.83–210.50 m CSF-A in Hole U1540D and 165.33–213.3 m CSF-A in Hole U1540E). This unit is marked by a significant increase in the proportion of calcareous Lithofacies 3 and 4 and the appearance of the clay-bearing to clay-rich Lithofacies 6 in the lowermost ~40 m of the unit. Lithofacies 7 occupies parts of the lowermost ~8 m of Unit II, is pale yellowish brown to dark reddish brown, and is visually enriched in iron (hydr)oxides. The last two sections of Core 383-U1540D-21H and the core catcher of Core 22H contain altered volcanic glass.

Biostratigraphic age assignments are overall consistent with each other and are mainly derived from diatom, radiolarian, and calcareous nannofossil datums as well as planktonic foraminifer biozonations. None of the different biozonations record any major hiatuses, indicating that a continuous sediment sequence from the early Pliocene through the Holocene was recovered at this site. The core catcher sample of the deepest core in Hole U1540E (Core 383-U1540E-10H) contained a few reworked taxa from the Miocene. However, the in situ microfossils present in this interval indicate a maximum early Pliocene age at the base of Site U1540.

Paleomagnetic measurements indicate a number of polarity changes that are best illustrated by downhole changes in inclination. The boundaries that define each polarity reversal are overall reasonably distinct. Prior to demagnetization, inclinations in the uppermost 40 m are dominantly positive, which is opposite to the expected inclination during the normal polarity Brunhes Chron. After demagnetization, inclinations become steeply negative and largely consistent with values expected for a geocentric axial dipole for the site's location. Positive inclinations associated with reversed polarity of the Matuyama Chron were observed in Holes U1540A, U1540B, and U1540D. However, the Matuyama–Brunhes polarity transition is not definitively captured within any core other than possibly at the base of Core 383-U1540B-5H. Three normal polarity subchrons within the Matuyama Chron were observed. The upper polarity transition of the Jaramillo Subchron was observed at the base of Core 383-U1540D-8H and somewhat more noisily in Core 383-U1540A-6H, and the lower Jaramillo polarity transition from the Matuyama Chron was observed in Cores 383-U1540B-10H and 383-U1540D-10H. The upper polarity transition from the Olduvai Subchron (C2n) was observed in Core 383-U1540B-17H and somewhat more noisily in Core 383-U1540A-12H. The lower polarity transition from the Matuyama Chron was observed in Core 383-U1540B-16H and possibly the upper part of Core 383-U1540A-13H. The normal polarity Reunion Subchron is recognized in all cored holes with varying fidelity. Deeper than ~180 m CCSF-A, polarity becomes more difficult to interpret as well-defined intervals of both polarities were observed on a decimeter scale.

We analyzed samples for headspace gas, interstitial water chemistry, and bulk sediment chemistry at a resolution of three samples per core from 0 to 120 m CSF-A, two per core from 120 to 150 m CSF-A, and one per core from 150 m CSF-A to the bottom of the

hole (Cores 383-U1540A-1H through 16H, 383-U1540B-2H, and 383-U1540D-15H through 21H). Methane concentrations are very low at this site overall, ranging from a maximum 6 ppm and a minimum of 2 ppm, with an average content of 4 ppm. Ethane and propane remained below detection limit throughout the entire hole.

Shallower than ~134 m CSF-A (Holes U1540A and U1540B), alkalinity and pH show no obvious trends, varying around averages of 3.4 ± 0.5 mM and 7.96 ± 0.07 , respectively. Below this depth, alkalinity and pH exhibit almost identical behavior, first increasing to a local peak at ~157 m CSF-A and then decreasing steadily to the bottom of Hole U1540D.

The composite carbonate record at Site U1540 shows downhole variability ranging from 6 to 93.5 wt% and a strong correlation with RGB blue and reflectance L^* (Figure F24) and GRA bulk density measurements. TOC content has a mean value of 0.36 wt% and a standard deviation of 0.13 across the entire cored interval. TOC values range from a maximum of 0.75 wt% at 20.63 m CSF-A in Hole U1540A, and a minimum of 0.13 wt% at 204.4 m CSF-A in Hole U1540D. The TOC record shows no correlation to the CaCO_3 record ($R^2 = 0.004$) downhole.

Physical property measurements at Site U1540 include nondestructive whole-round measurements of GRA bulk density, MS, WRMSL P -wave velocity, and NGR on core sections from Holes U1540A–U1540E. Additional physical properties collected include thermal conductivity on whole-round core from all holes. Data acquired from whole-round measurements are generally in good agreement with those from split-core measurements and discrete samples. In particular, discrete moisture and density (MAD) and GRA bulk densities correlate well, and GRA bulk density is also well correlated to discrete carbonate measurements from the different holes. Major bulk density peaks, which correspond to the lowest P -wave velocities, correlate with high-carbonate nannofossil content, whereas smaller GRA bulk density, NGR, and MS peaks observed simultaneously are partly linked to IRD content. Low GRA densities reflect the more biosiliceous facies and correspond to higher P -wave velocity and high porosity. The older part of the record at the site, from 185 m CCSF to the bottom, shows a clear inverse correlation of MS and GRA bulk densities, indicating that the oldest sediments are mainly composed of only two components, carbonate nannofossils and clay. NGR data also imply that total counts and the derived potassium content can be used as a semiquantitative proxy for the abundance of terrigenous material delivered by dust or sea ice and iceberg transport. Downhole changes in physical property characteristics overall are thus in good agreement with the identified lithostratigraphic units based on sedimentologic characteristics (Figure F25). Comparison of the different physical properties allows the discrimination of intervals consisting of mainly two components from intervals consisting of three or more components and could help provide a more quantitative determination of the sediment composition.

Correlations between holes were accomplished using the Correlator software (version 3.0). Tie points were established mostly using the RGB blue channel extracted from the Section Half Imaging Logger (SHIL) digital images, but in many cases a combination of measurements was used. We constructed a splice from 0 to 222.95 m CCSF-A using four holes (U1540A, U1540B, U1540D, and U1540E). However, because some misfired cores that were disturbed could not be used, the splice contains two gaps. The deepest cores of Holes U1540D and U1540E, 383-U1540D-22H and 23X and 383-U1540E-10H, were excluded from the spliced interval because of poor recovery and high levels of drilling disturbance.

We constructed a preliminary age model based on biostratigraphic and paleomagnetic age markers. These data suggest that the sedimentary sequence recovered at Site U1540 covers the past ~4.8 My (Figure F26). Though substantially lower than at Site U1539, sedimentation rates at Site U1540 are remarkably high for a pelagic setting. Below a condensed shallow section downhole to ~0.3 Ma, sedimentation rates average 7.5 cm/ky downhole to ~2.7 Ma and decrease to ~1.5 cm/ky during most of the Pliocene (2.7–4.8 Ma). This age model is generally consistent with preliminary stratigraphic tuning performed on board based on physical property data such as color measurements (RGB blue) and GRA bulk density.

The combination of nearly continuous recovery, very high sedimentation rates driven by high diatom productivity during glacial periods and potential sediment focusing, clear patterns in physical properties and sediment color, and a rich array of well-preserved diatoms combined with calcareous microfossils will provide unprecedented opportunities for improving our understanding of the dynamics of the ACC and its link to global carbon cycle changes at orbital and suborbital timescales.

Site U1541

Background and objectives

Site U1541 (Proposed Site CSP-1A) is located in the central South Pacific at $54^\circ 12.756'S$, $125^\circ 25.540'W$ ~1830 nmi west of the Magellan Strait at 3604 m water depth (Figure F27). The site sits at the western flank of the southernmost EPR, ~50 nmi north of the Tharp Fracture Zone and ~160 nmi west of the modern seafloor spreading axis, and is underlain by oceanic crust formed at the EPR at ~6–8 Ma (Eagles, 2006). Assuming overall constant seafloor half spreading rates of ~4.5 cm (Pitman and Heirtzler, 1966), the plate tectonic backtrack path of Site U1541 moves the site eastward, placing it ~100 nmi closer to the crest of the EPR at a water depth shallower by several hundred meters in the early Pliocene. At a smaller scale, the site is located in a ~4 nmi wide trough oriented NNE–SSW, parallel to the orientation of the EPR. The adjacent ridges rise to ~3400 m water depth southeast of the site and to ~3500 m water depth to the northwest (Figure F27).

Site U1541 is located on MCS Line AWI-201000014 at the intersection with Line AWI-201000016 (Figure F28A) (Gersonde, 2011). The seismic cross-lines indicate ~180 m thick sediments above oceanic basement. Sediments are mostly well stratified with flat lying, slightly irregular reflectors. Low to moderately reflecting layers become stronger deeper than ~120–140 m CSF-A. Sediment echo sound (Parasound) profiles (Gersonde, 2011) indicate moderate penetration (~80 m) with distinct layering (Figure F28B), suggesting a succession of fine-grained soft sediments with varying lithologic composition.

Site U1541 lies in the pathway of the SAP ACC, ~100 nmi north of the modern mean position of the SAF. In this sector of the ACC, the associated fronts are strongly steered by the topography of seafloor spreading systems (Udintsev and Eltanin-Tharp Fracture Zone systems).

SSTs seasonally vary between ~3°C (July–September) and ~7°C (January–March). The area is located west of the main Antarctic Intermediate and Mode Water formation regions in the Southeast Pacific. The 3604 m water depth places Site U1541 within LCDW (Figure F14).

The main objectives at Site U1541 were to

- Recover a moderate-resolution Subantarctic Miocene–Quaternary sediment record close to the SAF;

- Investigate the sequencing of calcareous and siliceous oozes, allowing for a wide range of paleoceanographic reconstructions;
- Reconstruct high-amplitude Subantarctic SSTs based on multiple proxies;
- Investigate long-term changes in dust input;
- Provide a record of LCDW and potential influence of AABW during glacial periods;
- Reconstruct productivity (opal versus carbonate), nutrient distribution, and dust-productivity coupling; and
- Recover a potential far-field record of WAIS variability.

Operations

During the transit to Site U1541, rough weather systems moved in, causing high seas at the site location. Therefore, upon reaching the site coordinates at ~2300 h on 15 June 2019, the vessel continued at reduced speed on a 245° weather avoidance course for another ~100 nmi past the site. At 2105 h on 16 June, the sea state had sufficiently calmed down to turn the ship around and head back.

We completed the sea voyage and arrived at Site U1541 at 0736 h on 17 June. We lowered the thrusters and the vessel was placed over the site coordinates and on dynamic positioning mode by 0820 h. Soon after, the drill crew started to build the APC/XCB BHA and to lower the drill string to the seafloor.

By 1815 h on June 17 2019, the drill bit was lowered to 3597 mbrf, 5 m above the seafloor depth calculated by the onboard PDR. We attempted to spud Hole U1541A, but the first core came up empty. Three more attempts followed with the bit at 3602, 3607, and 3617 mbrf before we successfully recovered any sediment. Finally, we spudded Hole U1541A at 2330 h on 17 June. Core 383-U1541A-1H recovered 9.5 m of sediment, indicating a missed mudline, and we ended the hole at that point.

The bit was raised to 3612.0 mbrf, and Hole U1541B was spudded at 0055 h on 18 June. Based on the recovery in Core 383-U1541B-1H, the seafloor was calculated at 3614.4 mbrf (or 3603.7 mbsl). Coring continued without incident until a partial stroke was recorded on Core 15H. When the core was recovered, the temperature tool (APCT-3) cutting shoe showed damage, and the driller pulled back the next APC core barrel before it arrived at the bottom. An XCB core barrel was dropped to attempt Core 16X. After drilling for 45 min and advancing only 0.5 m, the barrel was pulled back on board. Core 16X retrieved three large pebble-sized basalt pieces (2–5 cm in diameter) in the core catcher. The bit was pulled out of the hole, clearing the seafloor at 0100 h and ending Hole U1541B.

A total of 16 cores were taken in Hole U1541B. The APC system was used for 15 of these, reaching a depth of 138.0 m CSF-A before APC refusal and recovering 129.3 m (94%). The XCB system was used for one core; it advanced 0.5 m and recovered 0.06 m of basalt (12%). The APCT-3 was used on Cores 383-U1541B-4H, 7H, 11H, and 15H. Misfires were recorded on Cores 1H, 4H, and 6H, and a partial stroke was registered on Core 383-U1541B-15H. Twenty-five hours were spent in Hole U1541B.

The vessel was offset 20 m east of Hole U1541B, the bit was raised to 3608.0 mbrf, and Hole U1541C was spudded at 0325 h on 19 June. The hole was deepened with the APC system to 118.1 m CSF-A (Cores 383-U1541B-1H–13H) before coring was terminated at 1805 h on 19 June to allow the vessel to evade approaching heavy weather.

The drill pipe was retrieved, and the rig floor was secured for transit at 0204 h on 20 June, ending Hole U1541C and Site U1541. A total of 13 APC cores were taken over a 118.1 m interval with a recovery of 100.4 m (85%).

In summary, three holes were cored at Site U1541 using the full-length APC coring system. Hole U1541A was cored to 9.5 m CSF-A (9.67 m recovered; 101.8%), Hole U1541B was cored to 138.5 m CSF-A (129.27 m recovered; 93.3%), and Hole U1541C was cored to 118.1 m CSF-A (100.37 m recovered; 85%).

Principal results

A ~145 m spliced sedimentary sequence of Holocene–Miocene age was recovered at Site U1541 from Holes U1541A–U1541C (Figure F29). The sedimentary sequence includes four lithofacies, namely carbonate-bearing to carbonate-rich diatom ooze (Lithofacies 2), diatom-bearing to diatom-rich nannofossil/calcareous ooze (Lithofacies 3), nearly pure nannofossil ooze (Lithofacies 4), and clay-bearing to clayey biogenic ooze (Lithofacies 6) (Figure F30). Based on the distribution and co-occurrence of the defined lithofacies, we divided the Site U1541 sedimentary sequence into three lithostratigraphic units. The youngest, Unit I, is divided into two subunits: IA downhole to ~25 m CSF-A (~27 m CCSF-A) and IB from ~25 to ~63 m CSF-A (~66 m CCSF-A). The subunit boundary is determined by a transition from frequent and marked changes between Lithofacies 2, 3, and 4 to a dominance of Lithofacies 3. Unit II spans from ~63 to ~95 m CSF-A (~66 to ~98 m CCSF-A) and differs from Unit I above by an absence of diatom ooze (Lithofacies 2), resulting in a dominance of different varieties of nannofossil ooze (i.e., Lithofacies 3 and 4). Subunit IIIA, from ~95 to ~119 m CSF-A (~99 to ~127 m CCSF-A), can be distinguished from Unit II above by a distinct color change in Lithofacies 3 and 4 from white and light greenish gray to very pale brown and pale orange yellow and an increased contribution from Lithofacies 3. Subunit IIIB spans from ~119 to ~137 m CSF-A (~127 to ~145 m CCSF-A) and shows a dominance of Lithofacies 4 in contrast to Subunit IIIA. The presence of basalt rock and volcanic glass fragments at the base of Hole U1541B in Sections 15H-5 and 16X-CC suggests the complete recovery of the pelagic sediment sequence above oceanic basement at this site.

All microfossil groups studied on board (diatoms, radiolarians, silicoflagellates, benthic and planktonic foraminifers, nannofossils, and ostracods) are present at the site. The preservation of diatoms is moderate overall. Radiolarians are generally very well preserved except for a few samples. Silicoflagellates are rare to barren throughout the core except for four samples in which they are common to abundant. Calcareous nannofossils are present in all the samples. They are common to abundant except for a few samples and become dominant in specific nannofossil ooze intervals. Planktonic and benthic foraminifers and ostracods show good to moderate preservation in the upper part of Hole U1541B and moderate preservation in the lowermost core catcher samples. Ostracods are particularly abundant in Sample 9H-CC.

The biostratigraphic age model at Site U1541 was mainly based on analyses from Hole U1541B with additional samples analyzed for diatoms and nannofossils in Hole U1541C. Split-core samples were taken in both holes to refine the stratigraphy in the lower half of the site. In total, 117 biostratigraphic events were recognized and indicate an estimated age of ~7.7 Ma at the bottom of both holes. The integrated biostratigraphy of diatoms, radiolarians, calcareous nannofossils, and planktonic foraminifers showed no major hiatuses at Site U1541, suggesting a continuous sequence of late Miocene to Holocene sediments.

Paleomagnetic measurements indicate a number of polarity changes that are best illustrated by downhole changes in inclination. These capture a series of apparent polarity reversals that are consis-

tent between Holes U1541B and U1541C where they overlap and together provide a long and almost continuous record of polarity change over the past 8 My (Figure F31). The Matuyama/Brunhes and upper and lower Jaramillo polarity boundaries that define the upper part of the Matuyama Chron (C1r.1r and .1n) were observed. Most reversal boundaries from the Olduvai Chron (C2n) to C4r.1n (8.254 Ma) are generally well characterized. The record of inclination and intensity after 20 mT peak AF demagnetization for the splice composite record allows polarity interpretations and correlation to the geomagnetic polarity timescale (GPTS; Cande and Kent, 1995). Only the boundaries associated with the reversed polarity intervals of the Gauss Chron (C2An) (Kaena and Mountain) and the short normal polarity Subchron C3BR.2n that spans from 7.454 to 7.485 Ma are not clearly recognized over the 8.5 Ma interval.

We analyzed samples for headspace gas, interstitial water chemistry, and bulk sediment chemistry at a resolution of three per core from 0 to 9.5 m CSF-A in Hole U1541A and 0 to 138.5 m CSF-A in Hole U1541B and two per core from 0 to 32.6 m CSF-A in Hole U1541C. Methane concentrations are low at this site overall, never exceeding 3.47 ppmv. Ethane and propane remain below detection limit throughout the entire hole.

Alkalinity and pH decrease with depth starting around 80 m CSF-A. Interstitial water iron concentration deeper than 86 m CSF-A drops below detection limits, and interstitial water manganese concentration drops below detection limits deeper than 105.05 m CSF-A.

Calcium carbonate content is relatively high at this site; it never falls below 21.6 wt% and has a downhole average content of 68.6 wt%. Calcium carbonate content shows a good correlation with sediment color reflectance parameter L^* and RGB blue data (Figure F32). TOC contributes a maximum of 0.8 wt% to the total carbon pool throughout this site and shows no clear correlation with calcium carbonate content. TN is very low at this site, never exceeding 0.06 wt%.

The bulk sediment elemental composition shows marked increase in concentrations toward the bottom at 135.85 m CSF-A, indicating interaction between basement rocks and the sediment. Oxide contents of magnesium, potassium, iron, silica, and titanium co-vary strongly with aluminum oxides, suggesting that these elements are present primarily in clay minerals. High coherency between the downhole calcium and strontium concentration patterns suggests that strontium is incorporated in biogenic carbonate.

Physical property data acquired from whole-round measurements are generally in good agreement with those from split-core measurements for MS loop and point measurements. GRA bulk density and *P*-wave logger (PWL) velocity values indicate shorter term variations in Lithostratigraphic Unit I and the top of Unit II, and the shortest term variations at the top of Unit II and in Subunit IB are probably linked to 40 ky cycles in the early Pleistocene. MS shows both high amplitude and large timescale variability all along the record. GRA bulk density shows higher values in Unit II and Subunit IIIB, corresponding to a higher fraction of carbonate/nannofossils compared to diatoms, whereas MS and NGR show higher values in Subunit IIIA likely linked to higher terrigenous content.

Downhole changes in physical property characteristics overall are in good agreement with the defined lithofacies (Figure F33) but show different relationships with one another compared to Sites U1539 and U1540 in intervals of low diatom content.

Correlations between holes at Site U1541 were accomplished using Correlator software (version 3.0). Tie points were established

mostly using the core image RGB blue channel, but in many cases a combination of measurements was used. We constructed a splice from 0 to 127.254 m CCSF-A using three holes; however, because of some disturbed cores or cores that were too short to be used, the splice contains two gaps. Because Hole U1541B was drilled deeper than either of the other holes, the last two cores in Hole U1541B were appended to the bottom of the splice.

We constructed a preliminary age model based on biostratigraphic and paleomagnetic age markers. These data suggest that the sedimentary sequence recovered at Site U1541 covers the past ~8 My (Figure F34). Though substantially lower than at Site U1539 and U1540, sedimentation rates at Site U1541 average 2.7 cm/ky downhole to ~3 Ma and decrease to ~1.3 cm/ky during most of the Pliocene and late Miocene. This age model is generally consistent with preliminary stratigraphic tuning performed on board based on physical property data such as color measurements (RGB blue) and GRA bulk density.

The combination of nearly continuous recovery, medium high sedimentation rates primarily driven by calcareous and diatom productivity during glacial periods, clear patterns in physical properties and sediment color, and a rich array of well-preserved calcareous microfossils combined with diatoms will provide unprecedented opportunities for improving our understanding of the dynamics of the ACC and its link to global carbon cycle changes at orbital timescales back into the late Miocene. Site U1541 will therefore provide a crucial Miocene-to-present perspective on the evolution of the ACC system and its climate impact and will complement higher resolution paleoclimate reconstructions based on Sites U1539 and U1540 that cover the period from the Pliocene to present.

Site U1542

Background and objectives

Site U1542 (Proposed Site CHI-4B) is located at the Chile continental margin in the eastern South Pacific at 52°42.2917'S, 75°35.76915'W ~30 nmi west of the entrance of the Magellan Strait at 1101 m water depth (Figure F35). North of ~53°S, subduction in the Chile Trench is more orthogonal and results in a steep slope incised by canyons that run perpendicular to the trench axis. Recent sediment cover is minor at the lower slope, but a relatively small-scale sediment depocenter ("sediment drift") has been identified at the upper continental slope where Site U1542 is located.

Site U1542 is located on MCS Line AWI-20160511 ~1 nmi north of the intersection with Line AWI-20160513 (Figure F36A) (Lamy, 2016). The seismic cross-lines indicate sediments as thick as ~700 m above continental basement. The basement age of the drift is unknown; a maximum age of ~10 Ma can be derived from the end of the tectonic erosion and start of subsidence at the continental margin (Polonia et al., 2007). Sediments are mostly well stratified with slightly irregular reflectors. Low to moderately reflecting layers become stronger deeper than ~350 m sediment depth. Sediment echo sound (Parasound) profiles (Lamy, 2016) reveal moderate penetration (~50 m) with distinct layering (Figure F36B), suggesting a succession of fine- to medium-grained sediments with varying lithologic composition.

Site U1542 is located underneath the southward-flowing CHC, a northern branch of the ACC that continues toward the Drake Passage and provides a major fraction of the present-day northern Drake Passage transport (Well and Roether, 2003). Satellite-tracked surface drifters reveal that after crossing the EPR, Subantarctic surface water of the ACC is transported northeastward across the Southeast Pacific toward the Chile coast at ~45°S, 75°W (Figures F2,

F37). Here, presently only a minor part of ACC water is deflected northward into the HCS, whereas the major fraction is deviated southward toward the Drake Passage. The CHC thus transports a significant amount of northern ACC water toward the Drake Passage in a narrow ~100–150 km wide belt along the coast (Chaigneau and Pizarro, 2005) (Figures **F2**, **F37**). Modern surface current velocities in the CHC reach >35 cm/s, and high flow speeds of ~20 cm/s extend to mid-depths (Boisvert, 1969; Chaigneau and Pizarro, 2005).

Site U1542 is located at the lower limit of AAIW and lies close to or within the major modern AAIW formation area in the Southeast Pacific (e.g., Bostock et al., 2013). CDW underlies AAIW in this region (Figure **F38**). The area off the Strait of Magellan is located ~5° latitude north of the present SAF. Modern mean annual SST in this area is ~8°C, and the seasonal range is ~3°C.

The main objectives at Site U1542 were to

- Recover Pleistocene paleoceanographic records over the past several glacial–interglacial cycles with suborbital-scale resolution;
- Reconstruct the strength of the CHC (Subantarctic ACC) before entering the Drake Passage;
- Investigate AAIW and CDW water mass properties;
- Investigate changes of continental paleoclimate; and
- Recover a potential near-field record of Patagonian ice sheet variability.

Operations

Four holes were cored at Site U1542 using the full-length APC coring system: Holes U1542A (0–169.5 m CSF-A; 181.74 m recovered; 107%), U1542B (0–1.4 m CSF-A; 1.46 m recovered; 104%), U1542C (0–213.7 m CSF-A; 225 m cored and 236.9 m recovered; 105%), and U1542D (0–234 m CSF-A; 202.7 m cored and 205.4 m recovered; 101%).

Transit to Site U1542

At 0236 h on 20 June 2019, the vessel got underway at full speed northeast of Site U1541 to avoid heavy weather headed toward the operational area. Taking advantage of the transit, we deployed the towed magnetometer. The vessel continued transiting on a northeasterly course for 1109 nmi over 4.7 days until 1845 h on 24 June when sea conditions allowed us to turn around and start heading to our operational area at the southwest Chile margin. During the northward transit on 21 June, we passed just 120 nmi east of Point Nemo, the most remote location on our planet.

At 1845 h on 24 June, we set a southeast course toward Proposed Site CHI-1C, located at the southernmost Chile margin close to the Pacific entrance of the Drake Passage. This provided the best ability to change course for Sites U1542 or U1543 if the weather improved over those sites first. We continued on this course until 1230 h on 28 June, covering 873 nmi, when a strong system of high winds and seas was predicted to begin moving across the transit path and into the Chile margin operational area, forcing us to stop at 48°44.0'S, 089°24.7'W. The thrusters were lowered to wait for the system to pass.

At 1300 h on 1 July 2019, the seas and weather subsided enough over the operational area in the Chile margin to get underway again. The thrusters were raised, and the vessel was underway at 1330 h toward Site U1544. At 1345 h on 2 July, we decided to adjust the course for Site U1542.

Hole U1542A

We arrived at Site U1542 at 1948 h on 3 July 2019. The thrusters were lowered, and the vessel switched into dynamic positioning mode by 2030 h. An APC/XCB BHA was made up and lowered to the seafloor. Based on a PDR reading of 1111.4 mbrf, an APC shooting depth of 1108.4 mbrf was chosen, and Hole U1542A was spudded at 0215 h on 4 July. Core 383-U1542A-1H retrieved 7.1 m of sediment, determining the seafloor depth at 1110.8 mbrf (or 1099.8 mbsl).

APC coring continued until 165.6 m CSF-A (Cores 383-U1542A-1H through 19H) when the core barrel became detached from the sinker bars during retrieval. The core barrel fell ~50 m back to the landing seat. Three attempts were needed to finally retrieve the barrel. Once at the surface, we found that only 1.05 m of core was recovered, and we decided to switch to the HLAPC system to deepen the hole. The HLAPC system was used for one core, recording a partial stroke and recovering 3.91 m before the hole was abandoned because of high seas and winds. The bit was pulled to 1033 mbrf, clearing the seafloor at 2359 h on 4 July and ending Hole U1542A. A total of 20 APC and HLAPC cores were taken over a 169.5 m interval, recovering 181.74 m (107.2%).

Nonmagnetic core barrels and the core orientation tool were used on all cores. Formation temperature was measured on Cores 383-U1542A-4H, 7H, 10H, 13H, and 16H. Partial strokes were recorded on Cores 10H–13H and 16H–21F. The total time spent in Hole U1542A was 27.5 h (1.1 days).

Hole U1542B

The vessel waited on the seas and winds to calm from 0030 h to 1545 h on 5 July 2019 with the bit set at 1033 mbrf. Then, the vessel was offset 20 m east of Hole U1542A, the bit was lowered to 1104.0 mbrf, and Hole U1542B was spudded at 1745 h on 5 July. Core 383-U1542B-1H recovered only 1.46 m of sediment, and because of the low recovery and quality of the core we decided to obtain a better mudline core, so Hole U1542B was abandoned.

Hole U1542C

The vessel stayed at the same coordinates, and the bit was lowered to 1105.0 mbrf. Hole U1542C was spudded at 1830 h on 5 July 2019. Based on the recovery from Core 383-U1542C-1H (3.3 m), the seafloor was calculated at 1111.2 mbrf (1100.2 mbsl). APC coring continued to 169.3 m CSF-A with Core 20H. While running in with the core barrel for Core 21H, the Captain terminated operations because of high currents and winds. When the core barrel was retrieved, it was found that the pins had sheared and 3.57 m of core had been retrieved for an advance of 3.0 m.

At 0900 h on 6 July, the bit was pulled to 71.8 m CSF-A and the top drive set back to wait on weather with the bit in the hole. The crew continued to wait on weather until the seas subsided at 1630 h. The top drive was picked up, the hole was washed to the bottom by 1815 h, and coring continued with Cores 383-U1542C-22H–28H to 226.0 m CSF-A at midnight on 6 July.

A total of 26 cores were taken over a 217.0 m interval with two 4.5 m drilled intervals (9 m total drilled). A total of 228.91 m was recovered (106% recovery). Nonmagnetic core barrels were used with all cores, and the core orientation tool was removed after Core 383-U1542C-20H. A misfire was recorded on Core 8H, and partial strokes were recorded on Cores 10H, 12H, 13H, 16H, 17H, and 20H–28H.

Hole U1542D

The vessel was offset 20 m south of Hole U1542C, and the bit was spaced out to 1107 mbrf. Hole U1542D was spudded at 0600h on 7 July 2019. Based on recovery from Core 383-U1542D-1H, seafloor was calculated at 1111.7 m CSF-A (1100.7 mbrf). The hole then was cored to a total depth of 213.7 m CSF-A using the full-length APC system, and three intervals were drilled without recovery over a total of 11.0 m to help eliminate gaps in core coverage between holes. After reaching total depth, the bit was recovered to the surface, clearing the seafloor at 0245 h and the rotary table at 0545 h on 8 July. The BHA was racked back in the derrick, and the rig floor was secured for transit at 0555 h. The vessel was switched from dynamic positioning to cruise mode at 0558h, ending Hole U1542D and Site U1542.

Nonmagnetic core barrels were used on all cores, but no orientation or temperature measurements were taken. Partial strokes were recorded on Cores 383-U1542D-10H and 12H–27H. A total of 24 cores were taken over a 202.7 m interval with 205.4 m of recovery (101.2%). Three drilled intervals were used following Core 12H, 22H, and 24H. The hole was advanced by recovery on all partial strokes.

Principal results

A ~249 m spliced sedimentary sequence of Holocene–middle Pleistocene age was recovered at Site U1542 from Holes U1542A–U1542D (Figure F39). It comprises five lithofacies, of which one was identified at previous Expedition 383 sites in the central South Pacific (nannofossil ooze; Lithofacies 4). New lithofacies include clayey silt to silty clay (Lithofacies 8), carbonate- and/or sand-bearing clayey to sandy silt (Lithofacies 9), foraminifers, sand- and/or clay-bearing silty nannofossil ooze (Lithofacies 10), and sand (Lithofacies 11). Site U1542 is characterized by exceptionally high sedimentation rates that exceed 30 cm/ky. The entire recovered sedimentary sequence (~249 m) can be described as one lithostratigraphic unit (I) that covers the middle Pleistocene to Holocene interval. Sediments at Site U1542 are of siliciclastic nature. The sedimentary sequence at this site is dominated by dark gray to dark greenish gray clayey silts or silty clays, whereas thin beds of light greenish gray to light gray nannofossil ooze occur infrequently (Figure F40). Deeper than 50 m CSF-A, up to 3 m thick intervals of interbedded clayey silt and sand appear recurrently. In addition, thin, light gray, foraminifer-bearing calcareous ooze, clay-bearing silty nannofossil ooze, and foraminifer-rich nannofossil ooze was observed deeper than 95 m CSF-A.

Diatoms and radiolarians are rare throughout the sediment succession, silicoflagellates are absent, nannofossils are few to barren, benthic and planktonic foraminifers are abundant, and ostracods are sparsely present in some intervals at the site. The biostratigraphic age model for Site U1542 was mainly based on biostratigraphic analyses of core catcher and split-core samples from Hole U1542A (0–169 m CSF-A) and the bottom of Hole U1542C (172.3–234.0 m CSF-A). In total, eight biostratigraphic events were recognized that indicate an estimated age of ~0.7 Ma at the bottom of Hole U1542C.

The NRM of the archive-half sections of Site U1542 was measured and remeasured after alternating field (AF) demagnetization at 2 cm increments. In general, the number of demagnetization steps reflect

- The desire for more steps to study the magnetization(s),
- The severity of the drill string and natural overprints that hopefully can be removed to recover the natural remanence,

- The desire to use low peak fields to preserve the magnetization for future shore-based studies, and
- The need to maintain core flow through the laboratory.

After demagnetization, the intensity correlates with neither MS nor intensity prior to demagnetization, suggesting that the large drill string overprint was successfully removed and a geomagnetic signal was recovered. Inclinations are steep and positive prior to demagnetization and steep and negative after, suggesting, as with the intensity data, that a vertical (+z) drill string magnetic overprint was removed, giving way to a primary remanence that varies around expected directions (−69°) for the site latitude during normal polarity. The NRM after 15 mT shows no evidence for reversed polarity, suggesting that Site U1542 sediments are younger than 0.781 Ma (Hilgen et al., 2012). Shore-based studies will verify these interpretations and further develop these sediments as geomagnetic recorders, providing information on field intensity and directional variability while facilitating the next iteration in magnetic stratigraphy.

The interstitial water chemistry at this site is influenced by anaerobic organic matter remineralization. As oxygen is depleted with depth by respiration, nitrate, manganese, and iron serve as terminal electron acceptors in organic matter remineralization. Sulfate reduction occurs when other electron acceptors become consumed/unavailable, and once sulfate is completely consumed, methanogenesis becomes the primary remineralization pathway. At this site, sulfate reduction predominates in the shallow sediment and the sulfate–methane transition zone likely occurs at ~9 m CSF-A. This is not unusual for continental margin sediments and is probably due to the fact that Site U1542 lies in a high productivity zone supported by abundant supply of nutrients from continental runoff.

Calcium carbonate content is very low in this site with the exception of two peaks of 39.2 and 80.0 wt% at 97.22 and 125.57 m CSF-A, respectively. The carbonate record shows no correlation with reflectance L^* and RGB blue data. Organic carbon contributes a maximum of 0.67 wt% to the total carbon pool throughout this site. TN is very low, never exceeding 0.05 wt%, and low concentrations yield poor reproducibility of samples because of the detection limit of the instrument.

TOC/TN ranges between 0.82 and 18.56, suggesting a predominance of marine-derived organic matter. However, the very high TOC/TN values are likely artifacts due to the extremely low values and poor reproducibility of TN measurements.

Physical property data acquired from whole-round measurements for Site U1542 are generally in good agreement with those from split-core measurements; however, at this site considerable offsets in many individual section depths exist through all holes due to ongoing expansion of sediment between whole-round logging and splitting the cores. These differences will have to be corrected with further onshore measures such as high-resolution X-ray fluorescence scanning and manual inspection of MS parameters before sampling. The applied manual processing to all Site U1542 holes based on the whole-round WRMSL/Special Track Multisensor Logger (STMSL) log sheets provided sufficiently “clean” data for GRA bulk density, MS, and NGR data but yielded data loss of about 8%–15% depending on the hole and data category. *P*-wave velocity measurements largely failed for Site U1542 because of core expansion and related cracking of the sediment.

At this site, the downhole changes in physical property characteristics are only partly in agreement with the defined lithofacies (Figure F41) based on sedimentologic characteristics. The princi-

pally higher MS and GRA bulk density values match the hemipelagic composition and higher lithogenic input to the sedimentary matrix, but they show different relationships with one another compared to previous Sites U1539 and U1540 in intervals of low diatom content.

The higher complexity of this hemipelagic site in terms of sediment delivery mechanisms is reflected in the acquired physical properties of Site U1542. The predominantly terrigenous sediment supply likely is composed of multiple processes such as ice and glacial meltwater transport as well as dust and current-induced lateral material transport and sorting after deposition, all influenced by various forcing such as climate and glacier mass balance changes, associated sea level changes, and dynamic changes in mid-depth ocean circulation that affects the transport direction and strength on both orbital and short (sub-)millennial timescales.

We constructed a preliminary age model based on biostratigraphic age markers. These data suggest that the sedimentary sequence recovered at Site U1542 covers the past ~0.7 My (Figure F42) with sedimentation rates averaging 37 cm/ky. This age model is generally consistent with preliminary stratigraphic tuning performed on board based on physical property data such as color measurements (RGB blue) and GRA bulk density.

The combination of nearly continuous recovery, very high sedimentation rates in a sediment drift primarily driven by siliciclastic sediment input during glacial, and, though diluted, a rich array of calcareous microfossils combined with diatoms will provide unprecedented opportunities for improving our understanding of the dynamics of the northern ACC before entering the Drake Passage, variations of the CHC, intermediate water mass circulation, and Patagonian ice sheet variability.

Site U1543

Background and objectives

Site U1543 (Proposed Site ESP-1B) is located in the eastern South Pacific at 54°35.06'S, 76°40.59'W ~110 nmi west of the Chile coast and ~120 nmi south-southwest of Site U1542 at ~3860 m water depth (Figure F43). The site is situated west of the Chile Trench on a topographically elevated ridge ~300 m above the trench axis (~4200 m water depth) (Figure F44A). The basement at this location is oceanic crust older than ~20 Ma (Eagles, 2006).

Site U1543 lies at the intersection of two seismic lines with a sediment thickness of 600–700 m. MCS Line AWI-20160501 lies ~1 nmi southwest of the intersection with Line AWI-20160503 (Figure F44A) (Lamy, 2016). Sediments are mostly well stratified with only slightly irregular reflectors. Sediment cover is drape-type and entirely covers the underlying basement morphology (Figure F44A). Sediment echo sound (Parasound) profiles (Lamy, 2016) reveal moderate penetration (~70 m) with distinct layering (Figure F44B), suggesting a succession of fine-grained sediments with varying lithologic composition.

Site U1543 is located at the offshore reach of the southward-flowing CHC, a northern branch of the ACC that continues toward the Drake Passage and provides a major fraction of the present-day northern Drake Passage transport (Well and Roether, 2003). Satellite-tracked surface drifters reveal that after crossing the EPR, Subantarctic surface water of the ACC is transported northeastward across the Southeast Pacific toward the Chile coast at ~45°S, 75°W (Figures F2, F37). Here, presently only a minor part of ACC water is deflected northward into the HCS, and the major fraction is deviated southward toward the Drake Passage. The CHC thus transports a significant amount of northern ACC water toward the Drake

Passage in a narrow ~100–150 km wide belt along the coast (Chaigneau and Pizarro, 2005) (Figures F2, F37). Modern surface current velocities in the CHC reach >35 cm/s at the continental margin (Site U1542) but decrease offshore toward Site U1543 (Boisvert, 1969; Chaigneau and Pizarro, 2005).

Site U1543 is presently located at the lower limit of LCDW and might have been affected by AABW in the past (Figure F38). The site is located ~4° latitude north of the present SAF. Modern mean annual SST in this area is ~8.5°C, and the seasonal range is ~3°C.

The main objectives at Site U1543 were to

- Recover a Plio/Pleistocene undisturbed, continuous sediment record with medium high sedimentation rates;
- Reconstruct the strength of the CHC (Subantarctic ACC) before entering the Drake Passage;
- Investigate AABW, CDW, and Pacific Deep Water (PDW) water mass properties;
- Investigate changes of continental paleoclimate; and
- Recover a potential near-field record of Patagonian ice sheet variability over the past several million years.

Operations

The R/V *JOIDES Resolution* completed the 119 nmi transit from Site U1542 to Site U1543 at 1717 h on 8 July 2019, in 10.5 h and averaging 11.3 knots. A water depth reading using the Precision Depth Recorder set the preliminary seafloor depth at 3877.4 m below rig floor (mbrf) for Hole U1543A. The operations plan for Site U1543 initially included three holes to a depth of 400 m CSF-A. This was modified due to time lost to weather during the expedition, shallowing the penetration depth to 250 m CSF-A for each hole. While coring Hole U1543A, it was decided to continue deepening the hole until APC refusal, and to only core two holes to APC refusal at this site.

Hole U1543A

Hole U1543A was spudded at 0325 h on 9 July 2019, with Core 1H recovering 7.1 m. Core recovery determined a seafloor depth of 3863.4 mbsl. Soon after, at 0415 h, coring operations were suspended when one of the ship's main propulsion shafts was shut down due to mechanical problems. The engineers found the problem to be with a faulty shaft brake unit. The brake was taken out of service and the shaft was repaired and brought back online by 0900 h. At that point coring resumed and continued without interruption through 11 July in good weather and sea conditions. Coring was terminated after 90,000 lb of overpull were needed to retrieve Core 383-U1543A-36H from 339.6 m CSF-A. The drill string was pulled out of the hole, clearing the seafloor at 1830 h on 11 July. Overall, 36 cores were taken with 103% recovery.

Nonmagnetic core barrels were used on all cores. The core orientation tool was used in most cores except for 383-U1543A-8H, 14H, 16H, 17H, 24H–29H, and 32H–36H. Formation temperature measurements were taken with the APCT-3 on Cores 4H, 7H, 10H, 13H, 16H, and 19H. Partial strokes were recorded on Cores 32H–36H. The total time spent in Hole U1543A was 73.2 h (3.05 days).

Hole U1543B

The vessel was offset 20 m east of Hole U1543A, and Hole U1543B was spudded at 2010 h on 11 July 2019. The seafloor depth was calculated at 2876.4 mbrf (3865.3 mbsl) based on the recovery from Core 383-U1543B-1H. Hole U1543B was cored entirely with the APC system, and two drilled intervals were used to cover coring gaps between Holes U1543A and U1543B. The first drilled interval

(1.9 m) followed Core 1H, and the second drilled interval (1.5 m) followed Core 23H. Core 33H reached a total depth of 286.5 m CSF-A. Overall, 31 cores were taken over a 283.1 m interval with 104% recovery.

The bit was retrieved to the surface, clearing the seafloor at 1800 h on 13 July and the rotary table at 0045 h on 14 July. The rig floor was secured for transit at 0110 h, and *JOIDES Resolution* began transiting to Site U1544.

A total of 31 cores were taken over a 283.1 m interval. A total of 294.97 m was recovered (104.2% recovery). Two intervals were drilled without recovery for a total of 3.4 m. Nonmagnetic core barrels were used with all cores, and the core orientation tool was not used. Partial strokes were recorded on Cores 383-U1543B-31H-33H. The liners of Cores 3H, 14H, 20H, 21H, 25H, and 31H-33H shattered on the catwalk after holes were drilled to release formation gas.

Principal results

Site U1543 consists of a 376.3 m spliced sedimentary sequence (Figure F45, F46). The sedimentary record comprises five lithofacies (Figure F46) identified at previous Expedition 383 sites. These include major biogenic lithologies (nannofossil ooze [Lithofacies 4]; clay-bearing to clayey diatom ooze [Lithofacies 6]); and diatom-, clay-, and/or silt-bearing to rich nannofossil or calcareous ooze [Lithofacies 10] and mainly siliciclastic sediments (diatom-bearing to rich clay [Lithofacies 7] and silt-bearing clay to clay-bearing silt [Lithofacies 8]). The biogenic lithofacies were first characterized in the central South Pacific, whereas the pure siliciclastic sedimentary lithofacies were observed at the northern Chile margin Site U1542. Their definition is expanded according to the varieties of lithofacies observed at this site. The sedimentary sequence at Site U1543 is characterized by moderate average sedimentation rates of ~5 cm/ky. Site U1543 comprises two lithostratigraphic units, I (0–115 m CCSF-A) and II (115–376.3 m CCSF-A), that cover the early Pleistocene to Holocene and late Miocene to early Pleistocene intervals, respectively. Unit I largely consists of greenish gray to dark greenish gray silt-bearing clay (Lithofacies 8) that contains prominent 10–30 cm thick beds of light gray to light greenish gray carbonate-, clay-, and/or diatom-rich nannofossil ooze. Deeper than 115 m CCSF-A, the abundance of diatomaceous sediments increases, mostly in the form of diatom-bearing silty clay (Lithofacies 7) and carbonate and silt-bearing diatom ooze (Lithofacies 6). The latter often occurs in association with decimeter-scale beds of nannofossil ooze. Diatom mats and dropstones are rare, and burrows and mottling due to diagenetic overprints are frequent. Lithification, secondary carbonate precipitation, and microfracturing increasingly occur deeper than 275 m CCSF-A.

A ~339.54 m thick sediment succession of late Miocene age was recovered at Site U1543. Diatoms and radiolarians are abundant throughout the sediment succession, silicoflagellates are absent to few but locally abundant, nannofossil abundance varies strongly with good preservation, benthic and planktonic foraminifers are absent in glaciials and moderately preserved when present, and ostracods are rarely present at the site and restricted to Miocene samples. Biostratigraphic analyses of Hole U1543A core catcher and split-core samples allowed the recognition of 56 biostratigraphic events and indicate an estimated age of 7.3–8.5 Ma at the bottom of the hole.

The NRM of the archive-half sections of Site U1543 was measured and remeasured after AF demagnetization at 2 cm increments. In general, the number of demagnetization steps reflects

- The desire for more steps to study the magnetization(s),
- The severity of the drill string and natural overprints that hopefully can be removed to recover the natural remanence,
- The desire to use low peak fields to preserve the magnetization for future shore-based studies, and
- The need to maintain core flow through the laboratory.

The intensities before and after demagnetization for the upper 140 m of cores recovered from Hole U1543A are generally quite strong, varying around the 10^{-1} to 10^{-2} A/m range. A few discrete intervals have values as low as 10^{-4} A/m and are generally associated with MS lows. Deeper than 140 m CSF-A, intensities are slightly lower on average, falling to $\leq 10^{-2}$ A/m. Deeper than 240 m CSF-A, intensities are more variable and values in the 10^{-3} to 10^{-4} A/m range are much more common. Compared with previous sites, there is little difference in intensity before and after demagnetization at a peak AF of 15 mT, suggesting a finer magnetic mineralogy with a higher coercivity that is less susceptible to the drill string overprint.

The inclination for Holes U1543A and U1543B before and after AF demagnetization at 15 mT captures a series of apparent polarity reversals, either directly observed or inferred from polarity changes between cores, and when observed within a section the reversals are generally well defined and many times reproduced. Their positions in Hole U1543A are illustrated in Figure F47. All polarity boundaries from the Matuyama/Brunhes at 0.781 Ma through C3Br.2r at 7.285 Ma, 31 polarity reversals in total, are either directly observed or closely constrained and correlated to the GPTS (Cande and Kent, 1995; Hilgen et al., 2012). Reversed and normal polarity associated with the base of Chron C3B and the upper part of Chron C4 (older than 7.432 Ma) were observed, but the boundaries are not constrained in the deepest cores in Hole U1543A. Shore-based studies will refine correlations to the GPTS and facilitate development of relative paleointensity as the next steps in magnetic stratigraphy at Site U1543.

Biologically mediated anaerobic oxidation of organic matter is evidenced in the interstitial water by the presence of reduced iron and manganese, high alkalinity and pH, and high ammonium concentrations. Farther downcore, interstitial water chemistry may be additionally impacted by oxidative processes either deep in the core or below the cored depth with upward diffusion. This is demonstrated by decreases in ammonium and increases in sulfate with depth. The deeper parts of the core show evidence of carbonate dissolution as well as the interplay of dissolution and precipitation of iron- and manganese-bearing minerals.

Calcium carbonate content at this site is low but highly variable, and several peaks fluctuate between a minimum of 0.1 wt% and a maximum of 86.8 wt%. A linear correlation between calcium carbonate contents and RGB blue and reflectance L^* data was observed at this site for samples above 1 wt% (Figure F48). Organic carbon contributes a maximum of 0.43 wt% to the total carbon pool throughout this site.

TN is very low at this site, never exceeding 0.05 wt%. TOC/TN ranges between 0.34 and 19.91, suggesting a predominance of marine-derived organic matter. However, TOC/TN values closely mirror the TOC record, and TN values are likely too low to effectively apply TOC/TN values to sediments at Site U1543 (see discussion for Site U1542).

Data acquired from whole-round measurements for Site U1543 are generally in good agreement with those from split-core measurements. For Site U1543, MS, bulk density, and NGR are posi-

tively correlated with higher values that correspond to higher terrestrial component in the sediment (Lithofacies 7 and 8) and lower values that correspond to increased biogenic components (Lithofacies 4, 6, and 10). Although some minima in those physical properties corresponds to changing lithofacies, the physical properties also record subtle changes between terrestrial and biogenic components at high temporal frequency and for long term, allowing a precise reconstruction of warm and cold intervals and a precise correlation with isotopic stratigraphy. Furthermore, GRA bulk density and *P*-wave velocity show different behaviors for increased nanofossil and diatom presence in Lithofacies 8. Both values decrease for increasing nanofossil content, and GRA bulk density decreases. *P*-wave velocities, however, increase with increasing diatom content. It is thus possible, by combining the different high-resolution physical property records, GRA bulk density, magnetic susceptibility loop, and PWL, to interpret the large to smaller variations of the records as variations of either diatom or nanofossil content in terrigenous Lithofacies 8 (Figure F49).

We constructed a preliminary age model based on biostratigraphic and paleomagnetic age markers. These data suggest that the sedimentary sequence recovered at Site U1543 covers the past ~8.4 My (Figure F50). Sedimentation rates are overall relatively constant and average ~4.5 cm/ky. This age model is generally consistent with preliminary stratigraphic tuning performed on board based on physical property data such as color measurements (RGB blue) and GRA bulk density.

Overall, the undisturbed continuous sediment record with medium to high sedimentation rates extending into the Miocene recovered at Site U1543 is exceptional. The record is an important eastern end-member locality to the central South Pacific sites recovered during Expedition 383 that can be used to evaluate past oceanographic and climate changes from the late Miocene to the present. The observed variations in lithology and physical properties are at least in part driven by glacial–interglacial climate change that consistently impacts the delivery and nature of sediment deposited at Site U1543. The longer term evolution of these depositional processes recorded at Site U1543 will illuminate the multimillion-year history of Patagonian ice sheet growth and retreat, ACC transport, and the underlying climatic conditions in which these processes evolved.

Site U1544

Background and objectives

Site U1544 (Proposed Site CHI-1C) is located at the Chile continental margin in the eastern South Pacific at 55°32.2192'S, 71°35.6194'W ~30 nmi southwest off the Chile coast and at ~2090 m water depth (Figure F51).

Tectonically, the southern Chile margin is characterized by a complex geodynamic setting with oblique convergence between plates, transcurrent motion, and tectonic rotation on land (Polonia et al., 2007). The geodynamic setting of the southernmost Chile margin is driven by relative movements between three main plates: Antarctica, Scotia, and South America. This complex setting results in strong segmentation of the southern Chile margin as clearly visible in bathymetric data and multichannel seismic data (Polonia et al., 2007) (Figure F4). There are two major fore-arc basins at ~53°–54.5°S and 55°–56.5°S that are characterized by a thick sediment infill of up to ~3 km. Site U1544 is situated close to the axis of the southern fore-arc basin on the continental crust of the Scotia plate. This southern basin most likely formed after the ridge consumption (10–14 Ma). When accretion resumed, it contributed to an outer

high that provided the barrier for the accumulation of continent-derived sediments. The upper unit in the southern basement, relevant for drilling, shows onlap terminations on both basin flanks.

Site U1544 lies on multichannel seismic (MCS) Line IT95-181, ~3 nmi southeast of the intersection with Profile IT95-171 (Figure F52) (Lamy, 2016). Sediments are mostly well stratified and sediment cover at the site exceeds 2.5 km. Sediment echo sound (Parasound) profiles (Lamy, 2016) reveal poor to moderate penetration (~30 m) with distinct layering and an irregular surface, suggesting a succession of fine- to medium-grained sediments with occasional coarser grained composition.

Site U1544 is located underneath the southward-flowing CHC, a northern branch of the ACC that continues toward the Drake Passage and provides a major fraction of the present-day northern Drake Passage transport (Well and Roether, 2003). Satellite-tracked surface drifters reveal that after crossing the East Pacific Rise Subantarctic surface water of the ACC is transported northeastward across the Southeast Pacific toward the Chile coast at ~45°S, 75°W (Figures F2, F37). Here, presently only a minor part of ACC water is deflected northward into the HCS, whereas the major fraction is deviated southward toward the Drake Passage. The CHC thus transports a significant amount of northern ACC water toward the Drake Passage within a narrow belt of ~100–150 km width along the coast (Chaigneau and Pizarro, 2005) (Figures F2, F37). Modern surface current velocities in the CHC reach >35 cm/s, and high flow speeds of ~20 cm/s extend to mid-depths (Boisvert, 1969; Chaigneau and Pizarro, 2005).

Site U1544 is presently located at the lower limit of Lower Circumpolar Deep Water (LCDW) and might have been affected by AABW in the past (Figure F38). The site is located ~4° north of the present SAF. Modern mean annual SST in this area is ~8.5°C, and the seasonal range is 6°–9°C.

The main objectives at Site U1544 were to

- Recover Pleistocene paleoceanographic records over the past several glacial–interglacial cycles with suborbital-scale resolution,
- Reconstruct the strength of the CHC (Subantarctic ACC) before entering the Drake Passage,
- Investigate CDW and PDW water mass properties,
- Investigate changes of continental paleoclimate, and
- Recover a potential near-field record of Patagonian ice sheet variability.

Operations

We arrived at Site U1544 at 1746 h on 14 July 2019. Soon after arrival, the rig floor crew assembled the advanced piston corer/extended core barrel (APC/XCB) bottom-hole assembly (BHA) and deployed the drill string to the seafloor, reaching 2055 m below rig floor (mbrf) by midnight.

The initial attempt to spud Hole U1544A was made at 0015 h on 15 July and resulted in a bent/broken APC core barrel, which made it necessary to pull the bit back to the rig floor and remove the stuck portion of the barrel. The vessel was then moved 50 m southwest at 225° to attempt starting Hole U1544A again. Due to a hard layer below the sediment surface, we used the XCB system to spud the hole and break through the hard layer.

Hole U1544A finally was spudded at 1330 h on 15 July with the first core advancing 7.8 m into a softer formation. When the XCB core barrel was retrieved, we found a ~5 cm granite pebble in the core catcher. We then deployed the APC system and cored to APC

refusal at 88.1 m below seafloor (m CSF-A). The half-length APC (HLAPC) system was used to deepen the hole to the final depth of 106.0 m CSF-A before coring was terminated due to high seas and winds. The drill string was pulled out of the hole, clearing the seafloor at 0958 h on 16 July and ending Hole U1544A.

The vessel was offset 20 m to the east of Hole U1544A with the bit at 2055 mbrf to wait for the seas to subside, so an attempt could be made to spud Hole U1544B. However, heave was still above 3.5 m at 0630 h on 17 July and operations were terminated. The drill pipe was pulled to the surface and the vessel was secured for transit by 1300 h on 17 July, ending Site U1544. A total of 19 cores were taken over a 103.0 m interval with 89% recovery.

Principal results

Site U1544 consists of a 91.3 m sedimentary sequence obtained from a single hole (Figure F53). The sedimentary record comprises four lithofacies (Figure F54), all of which were identified at previous Expedition 383 sites. They include fine-grained (biosilica-bearing silty clay [Lithofacies 7] and silty clay [Lithofacies 8]), coarse-grained (sand [Lithofacies 11]) siliciclastic sediments, and biogenic sediments (diatom-, clay-, and/or silt-bearing to -rich nannofossil or calcareous ooze [Lithofacies 10]). The sedimentary sequence in Hole U1544A is dominated by siliciclastic sediments (i.e., Lithofacies 7, 8, and 11), with only sporadic layers of calcareous ooze. Sedimentation is characterized by highly variable sedimentation rates of ≥ 25 cm/ky, with rapidly deposited gravity-driven sedimentation (e.g., turbidities) contributing a significant fraction of the sedimentary sequence. Hole U1544A sediments make up only lithostratigraphic unit in this hole, Unit I (7.7–106.1 m CSF-A), which covers the late Pleistocene interval. Unit I largely consists of greenish gray to dark greenish gray silty clay (Lithofacies 8) or biosilica-bearing silty clay (Lithofacies 7) that frequently alternates with prominent decimeter- to meter-scale dark gray sand layers that often have normal grading and erosional boundaries to underlying strata. Prominent <1 m thick beds of light gray to light greenish gray silt-bearing and foraminifer-rich nannofossil ooze occur in two intervals at 26–28 and 80.5–82.5 m CSF-A. Dropstones are rare, whereas mottling due to diagenetic overprints is minor but frequent. The upper core, 383-U1544A-2X, retrieved a single hard rock pebble that suggests a rocky and hard substrate close to the seafloor at this site.

Diatoms are rare to abundant throughout the sediment succession, radiolarians and silicoflagellates are absent, nannofossils are abundant to barren, benthic and planktonic foraminifers are abundant above ~85 m CSF-A, and ostracods are sparsely present throughout the sequence, but abundant at ~60 and ~101 m CSF-A. The biostratigraphy of Site U1544 is not well constrained because of the lack of marker species for most fossil groups. Based on the biostratigraphic events recorded, the estimated age is <0.42 Ma at the bottom of Hole U1544A.

The NRM of the archive-half core sections was measured before and after AF demagnetization at 2 cm increments. The intensities before and after demagnetization for the upper 140 m of cores recovered from Hole U1544A are generally quite strong, varying around the 10^{-1} to 10^{-2} A/m range. A few discrete intervals have values as low as 10^{-4} A/m and are generally associated with magnetic susceptibility (MS) lows. Below 140 m CSF-A, intensities are slightly lower on average, falling in the 10^{-2} A/m and below range. Below 240 m CSF-A, intensities are more variable, with values in the 10^{-3} to 10^{-4} A/m range being much more common. Compared with previous sites, there is little difference in intensity before and after demagnetization at a peak field of 15 mT, suggesting a finer magnetic

mineralogy with a higher coercivity that is less susceptible to the drill string overprint. Although a very noisy record, no evidence for reversed polarity is observed, suggesting that Site U1544 sediments are younger than 0.781 Ma (Hilgen et al., 2012).

We constructed a preliminary age model based on biostratigraphic and paleomagnetic age markers. These data suggest that the sedimentary sequence recovered at Site U1544 covers less than ~0.4 Ma. Sedimentation rates are high and average ~25 cm/ky.

The low sampling resolution makes it difficult to interpret interstitial water profiles from this site. Low sulfate and high methane concentrations suggest that the sulfate–methane transition zone occurs shallower than ~10 m CSF-A (Iverson et al., 1985). The high alkalinity indicates that organic matter degradation is dominated by anaerobic processes, and concentrations of interstitial water iron suggest that microbial iron reduction also occurs at this site. Because of limited time, only two samples were analyzed for TN and carbon, and observed TN values are among the highest of all Expedition 383 sites.

At this site, the downhole changes in physical property characteristics do not show obvious features aligned with the defined lithofacies based on sedimentologic characteristics (Figure F55). Principally, lower MS and low GRA bulk density values match the interval of higher biogenic component, but they do not show a unique relationship with one another throughout the record. Comparing NGR and MS records could provide indications about the different terrigenous components because they are mainly anticorrelated like at the shallow-water Site U1542.

The close proximity of Site U1544 to the southernmost Chile margin explains the dominance of terrigenous sediments at this site. Terrigenous sediment is likely delivered to the site by a combination of ice rafting, suspension fallout from glacial meltwater plumes and/or freshwater plumes that originate from nonglaciated catchments, episodic downslope transport from the outer continental shelf, and fine-grained sediments transported by the CHC entering the Drake Passage as the northern branch of the ACC. The large proportion of normally graded sand beds with erosive basal contacts implies that turbidity currents were a transport mechanism to carry sediments from the continental slope and the upper slope during the Middle and Upper Pleistocene. We conservatively estimate the contribution of turbidities as ~40% based on the thickness of sand layers (lithofacies 11). However, it is possible that a certain fraction of the hemipelagic sediments also represents a continuation of the normal graded turbidite deposition sequence; if so, then turbidites may account for a larger proportion of the sequence and sedimentation at Site U1544 was rather discontinuous. Future geochemical and sedimentary analyses will provide crucial insights into the nature and temporal evolution of sedimentation processes at Site U1544.

Preliminary scientific assessment

Overall assessment

The overall goal of Expedition 383 was to improve our knowledge of Plio/Pleistocene atmosphere–ocean dynamics of the ACC in the Pacific and their implications for regional and global climate and atmospheric CO₂ based on sediment records with highest possible resolution.

Expedition 383 was very successful. We accomplished drilling and coring at six of the planned seven sites in the South Pacific. The drilling and coring objectives were largely achieved, and a total of 2636 m of core was recovered from six sites over a 2649 m cored

interval. This is the first time sedimentary records of this length and, in particular, of this resolution have been recovered from the South Pacific. We achieved our overall goal of recovering sedimentary sequences to improve our knowledge of Plio/Pleistocene atmosphere-ocean dynamics. Based on extrapolating sedimentation rates from the site survey cores and planned length of the drilled sequences, we expected to reach back to the early Pliocene (~5 Ma). At two of our sites (U1541 and U1543), we were able to drill continuous sequences that reach back to about 8 Ma, well into the Miocene, thus exceeding our expectations and allowing an even more complete investigation of the long-term evolution of the climate system in the South Pacific. The key achievements of the drilling operations include the following:

- All sequences cored are continuous and do not display any hiatuses.
- Site U1539 and to a lesser extent Site U1540 have very high sedimentation rates for a pelagic setting, thus allowing for the first time analyses of climate variability at millennial and submillennial timescales in the Southern Ocean. The sites are complemented by a lower resolution site, U1541, with a remarkably constant sedimentation rate of 2.7 cm/ky downhole to ~3 Ma and then ~1.3 cm/ky during most of the Pliocene and late Miocene. It provides a continuous paleoclimate record back to the Miocene that is expected to serve as a reference site for the region.
- Sites U1541 and U1543 provide the longest continuous paleoceanographic records obtained in the Southern Ocean.
- Even though the cross-frontal transect in the central South Pacific could not be cored, Site U1539, closest to the modern SAF, and Site U1540, ~170 nmi north of the front, provide pronounced lithologic differences, suggesting strong gradients between both sites, particularly during glacials.
- The two long-timescale cores from the central South Pacific (Site U1541) and the eastern South Pacific (Site U1543) provide an excellent zonal transect across the eastern part of the South Pacific over the past 8 My.

Despite the drilling success at the six sites, the overall expedition experienced two challenges that affected the drilling strategy and, to some extent, the scientific objectives:

- Approval: By the time we left Punta Arenas on May 24, we had not obtained approval from the Chilean authorities to conduct drilling and coring operations in their territorial waters. This forced us to change the original operations plan by reordering the sequence of our sites to allow the Chilean authorities more time for approval for the initial sites planned in Chilean waters. We received notice about clearance on 31 May 2019, 1 week into the expedition when we were already well underway to the drill sites in the central South Pacific. The delay in obtaining the permits severely limited our flexibility in planning operations.
- Weather-related interruptions and adjustments: During the entire expedition, we faced rough weather and sea conditions that impacted our drilling operation and required us to adjust our drilling plans on a continuous basis. Weather-related challenges were anticipated because Expedition 383 was targeting areas of the South Pacific Ocean that are within the range of maximum wind speeds of the Southern Westerly Winds. Expedition 383 was scheduled during austral autumn/winter when historical weather records show winds speeds in the Pacific Southern Ocean to be substantially reduced. However, we encountered

rough weather and seas resulting in the loss of substantial operational time above the long-term climatological mean. Broadly, the weather-related issues were two different types:

- Weather and sea conditions on site that stopped coring operations and partly prohibited the coring of three full holes per site as originally planned. At most of the sites, we had to interrupt drilling operations because of high seas and resulting high heave. In these cases, we stayed on site, waited until the seas calmed sufficiently, and continued drilling. Such conditions lead to 106 h of lost time.
- While drilling at Site U1541 in the central South Pacific, a large-scale (>2000 km) storm system developed in the Southeast Pacific and forced us to transit on a northeast course across the South Pacific on a bad-weather avoidance course. This not only caused a significant loss of time (12.3 days) but also made it impossible to reach one of our targeted sites, Proposed Site CSP-3A, at 60°S close to the PF. As a result of the bad-weather avoidance, we had to eliminate Site CSP-3A from our planned drilling schedule.

Despite encountering weather-related challenges, we were successful in mitigating the impacts on the scientific drilling program through good time management, intense attention to weather forecasting, and excellent communication with the captain and the operations superintendent.

Assessment related to the scientific hypotheses

Expedition 383 was built around two major scientific hypotheses. Here, we break down the objectives of the expedition in the context of these two hypotheses and document the achievements and potential for each hypothesis.

Hypothesis H1: ACC dynamics and Drake Passage throughflow conditioned the global Meridional Overturning Circulation and high-low climate linkages on orbital and submillennial timescales since the Pliocene.

The Drake Passage is the major geographic constriction for the ACC and forms an important pathway for the return of upper waters to the Atlantic through the cold-water route of the MOC (Gordon, 1986). Resolving changes in the flow of circumpolar water masses through the Drake Passage is crucial for advancing our understanding of the Southern Ocean's role in affecting ocean and climate change on a global scale. Before Expedition 383, reconstructions of the throughflow along the Chile margin were limited to the past 60 ky, and no further information was available. We were highly successful in drilling Site U1542, which will extend this type of record back to the middle Pleistocene. Site U1542 is complemented by a long and detailed record from Site U1543 in the ESP that reaches back to the Miocene. The weather conditions at Site U1543 were extremely favorable for the region and allowed us to drill two holes deeper than 300 m without any interruption and a minimum of heave-related drilling disturbance. The record recovered at this site has excellent paleomagnetism and a pronounced variability in MS that provides a promising tuning target. At the third site in the region, U1544, the borehole depth had to be shortened to about 100 m because of rough weather and time constraints at the end of the expedition. Accordingly, the record at this site is much shorter and limited to the last few glacial cycles but will prove helpful to evaluate Hypothesis H1 with regard to the last interglacial period, MIS 5, and potentially MIS 7. However, two fully occupied sites (U1542 and U1543) provide continuous and very well dated se-

quences that include relatively highly resolved interglacials and glacial–interglacial transitions. The two sites together will allow us to address the long-term, orbital-scale, and millennial-scale evolution and variability in ACC strength and Drake Passage throughflow, particularly in warmer-than-present interglacials (e.g., MIS 5e, 11, and 31) and across the Pliocene warm period at Site U1543. We anticipate that the recovered material will allow us to fully tackle Hypothesis H1 and associated research questions, even exceeding the timescale to the late Miocene.

Hypothesis H2: Variations in the Pacific ACC determine the physical and biological characteristics of the oceanic carbon pump and atmospheric CO₂.

Atmosphere-ocean-cryosphere interactions and teleconnections between high and low latitudes play an important role in understanding processes and feedbacks of past and future climate change, and the Subantarctic Southern Ocean provides the major link between Antarctica and the low latitudes. In the Southern Ocean, these interactions are believed to control sea ice cover, AIS dynamics, upper ocean stratification, biological nutrient utilization, and exposure rates of deep water. They have been considered to play a key role in explaining the variability in atmospheric CO₂ concentrations.

We drilled three excellent sites in the Subantarctic South Pacific, U1539, U1540, and U1541, that matched or even exceeded our expectations for these records in resolution (Site U1539) and covered time period (Site U1541). These drill cores from the Pacific sector of the Southern Ocean will allow us to explore in more detail the glacial–interglacial changes and longer term climate evolution North of the Antarctic PF. Our drill sites from the central South Pacific (U1539, U1540, and U1541) will provide robust data to test to what extent processes that are active in the Atlantic sector can be translated to the Pacific sector, thus allowing the construction of a more global picture of the Southern Ocean’s role in nutrient distribution and biogenic export production and their impact on CO₂ variations. The cores will enable us to generate key records of SST, dust, export production, and other parameters back to the Miocene. These reconstructions will provide important benchmarks for a climate-sensitive region that are important constraints to more realistically model the impact on the ocean carbon cycle under different climate boundary conditions.

Unfortunately, we were not able to reach the site in the Polar Frontal Zone (Proposed Site CSP-3A) because of persistent rough weather in the region, and therefore we were not able to complete the cross-frontal transect. The elimination of this key site from Expedition 383 has impacts on some aspects of testing Hypotheses H2 and related research questions that target processes in the Antarctic zone south of the PF as well as comparisons between the Subantarctic and Antarctic zone.

At two of the sites (U1540 and U1541) east and west of the EPR, we drilled to the basement of the overlying sediment sequence. The lowermost sediments at both sites provide evidence of hydrothermal alteration that may add an unforeseen aspect to the future research at the Expedition 383 sites.

The potential range of scientific applications for the unique set of cores drilled during Expedition 383 is very large, encompassing paleoclimate reconstructions, carbon cycle, evaluation of hydrothermal inputs, geochemistry, reconstruction of ice sheet variability, land-ocean interaction, and sedimentary processes. We anticipate exciting and important results from both the postexpedi-

tion research and research endeavors beyond the science party for many years to come.

Outreach summary

Expedition 383 shipboard participants included an outreach officer from the United States. She communicated the goals of the expedition to students and the general public across the globe through live broadcasts and social media. She also developed a virtual field trip education outreach activity targeted toward high school and undergraduate students.

Live Zoom broadcasts

Using the direct satellite link, an iPad, and the Zoom videoconferencing software, the education outreach officer was able to help the expedition scientist connect with people all over the world. A typical session included an introduction to the current location and expedition goals followed by a tour of the laboratories, tracing the core path as it comes on board. Scientists were stationed at each laboratory to give an overview of the science being conducted and to answer questions. We finished each tour with a 10–15 min question and answer session. During Expedition 383, we completed 23 broadcasts and connected with 10 different countries, including 5 different states in the United States. We also worked with the National Science Foundation to hold a Facebook Live event that reached over 2700 views.

Social media

During the expedition, the education outreach officer managed the *JOIDES Resolution* Expedition 383 website (<https://joidesresolution.org/expedition/383/>) and completed a total of 35 Career Spotlight blogs that highlighted the careers of the expedition crew members. She assisted scientist Anieke Brombacher, who posted in both English and Dutch, with access to the website. Together they posted a total of 48 blogs. The expedition website also has a link to the Columbia University Earth Institute State of the Planet blog series where Co-Chief Gisela Winkler and scientists Julia Gottschalk and Jenny Middleton posted articles about Expedition 383 (<https://blogs.ei.columbia.edu/features/ocean-of-extremes/>).

The education outreach officer also added at least one post every day to the *JOIDES Resolution* Facebook, Twitter, and Instagram social media accounts. This resulted in an increase of 89 new Instagram followers, 173 new Twitter followers, and 415 new Facebook page likes over the 2 month expedition. That is a total increase of 677 newly engaged individuals.

Research documentary production

During Expedition 383, the *JOIDES Resolution* Science Operator (JRSO) granted Maylo Films, Inc., limited access to *JOIDES Resolution* to make recordings for a documentary motion picture. The recordings made by Brian Cimagala (Film Producer) and Alejandro Wilkins (Director of Photography) relate to the expedition’s scientific mission and chronicle the tasks and efforts of the scientists, the technicians and the ship’s crew who participated during Expedition 383.

References

- Anderson, R.F., Ali, S., Bradtmiller, L.I., Nielsen, S.H.H., Fleisher, M.Q., Anderson, B.E., and Burckle, L.H., 2009. Wind-driven upwelling in the

- Southern Ocean and the deglacial rise in atmospheric CO₂. *Science*, 323(5920):1443–1448. <https://doi.org/10.1126/science.1167441>
- Anderson, R.F., Barker, S., Fleisher, M., Gersonde, R., Goldstein, S.L., Kuhn, G., Mortyn, P.G., Pahnke, K., and Sachs, J.P., 2014. Biological response to millennial variability of dust and nutrient supply in the Subantarctic South Atlantic Ocean. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 372(219):20130054. <https://doi.org/10.1098/rsta.2013.0054>
- Barker, S., and Diz, P., 2014. Timing of the descent into the last ice age determined by the bipolar seesaw. *Paleoceanography and Paleoclimatology*, 29(6):489–507. <https://doi.org/10.1002/2014PA002623>
- Barrows, T.T., Juggins, S., De Deckker, P., Calvo, E., and Pelejero, C., 2007. Long-term sea surface temperature and climate change in the Australian–New Zealand region. *Paleoceanography and Paleoclimatology*, 22(2):PA2215. <https://doi.org/10.1029/2006PA001328>
- Beal, L.M., De Ruijter, W.P.M., Biastoch, A., Zahn, R., and SCOR/WCRP/IAPSO Working Group 136, 2011. On the role of the Agulhas system in ocean circulation and climate. *Nature*, 472(7344):429–436. <https://doi.org/10.1038/nature09983>
- Benz, V., Esper, O., Gersonde, R., Lamy, F., and Tiedemann, R., 2016. Last Glacial Maximum sea surface temperature and sea-ice extent in the Pacific sector of the Southern Ocean. *Quaternary Science Reviews*, 146:216–237. <https://doi.org/10.1016/j.quascirev.2016.06.006>
- Boisvert, W.E., 1969. *Technical Report: Major Currents off the West Coasts of North and South America*: Washington, D.C. (Naval Oceanographic Office). <https://catalog.hathitrust.org/Record/008348460>
- Bostock, H.C., Sutton, P.J., Williams, M.J.M., and Opdyke, B.N., 2013. Reviewing the circulation and mixing of Antarctic Intermediate Water in the South Pacific using evidence from geochemical tracers and Argo float trajectories. *Deep Sea Research, Part I: Oceanographic Research Papers*, 73:84–98. <https://doi.org/10.1016/j.dsr.2012.11.007>
- Bradt Miller, L.I., Anderson, R.F., Fleisher, M.Q., and Burckle, L.H., 2009. Comparing glacial and Holocene opal fluxes in the Pacific sector of the Southern Ocean. *Paleoceanography and Paleoclimatology*, 24(2):PA2214. <https://doi.org/10.1029/2008PA001693>
- Brierley, C.M., Fedorov, A.V., Liu, Z., Herbert, T.D., Lawrence, K.T., and LaRiviere, J.P., 2009. Greatly expanded tropical warm pool and weakened Hadley circulation in the early Pliocene. *Science*, 323(5922):1714–1718. <https://doi.org/10.1126/science.1167625>
- Brovkin, V., Ganopolski, A., Archer, D., and Rahmstorf, S., 2007. Lowering of glacial atmospheric CO₂ in response to changes in oceanic circulation and marine biogeochemistry. *Paleoceanography and Paleoclimatology*, 22(4):PA4202. <https://doi.org/10.1029/2006PA001380>
- Callahan, J.E., 1972. The structure and circulation of deep water in the Antarctic. *Deep-Sea Research and Oceanographic Abstracts*, 19(8):563–575. [https://doi.org/10.1016/0011-7471\(72\)90040-X](https://doi.org/10.1016/0011-7471(72)90040-X)
- Cande, S.C., and Kent, D.V., 1995. Revised calibration of the geomagnetic polarity timescale for the Late Cretaceous and Cenozoic. *Journal of Geophysical Research: Solid Earth*, 100(B4):6093–6095. <https://doi.org/10.1029/94JB03098>
- Caniupán, M., Lamy, F., Lange, C.B., Kaiser, J., Arz, H., Kilian, R., Baeza Urrea, O., et al., 2011. Millennial-scale sea surface temperature and Patagonian Ice Sheet changes off southernmost Chile (53°S) over the past ~60 kyr. *Paleoceanography and Paleoclimatology*, 26(3):PA3221. <https://doi.org/10.1029/2010PA002049>
- Chaigneau, A., and Pizarro, O., 2005. Surface circulation and fronts of the South Pacific Ocean, east of 120°W. *Geophysical Research Letters*, 32(8):L08605. <https://doi.org/10.1029/2004GL022070>
- Chalk, T.B., Hain, M.P., Foster, G.L., Rohling, E.J., Sexton, P.F., Badger, M.P.S., Cherry, S.G., et al., 2017. Causes of ice age intensification across the Mid-Pleistocene Transition. *Proceedings of the National Academy of Sciences of the United States of America*, 114(50):13114–13119. <https://doi.org/10.1073/pnas.1702143114>
- Chase, Z., Anderson, R.F., Fleisher, M.Q., and Kubik, P.W., 2003. Accumulation of biogenic and lithogenic material in the Pacific sector of the Southern Ocean during the past 40,000 years. *Deep Sea Research, Part II: Topical Studies in Oceanography*, 50(3–4):799–832. [https://doi.org/10.1016/S0967-0645\(02\)00595-7](https://doi.org/10.1016/S0967-0645(02)00595-7)
- Chase, Z., Kohfeld, K.E., and Matsumoto, K., 2015. Controls on biogenic silica burial in the Southern Ocean. *Global Biogeochemical Cycles*, 29(10):1599–1616. <https://doi.org/10.1002/2015GB005186>
- Cook, C.P., van de Flierdt, T., Williams, T., Hemming, S.R., Iwai, M., Kobayashi, M., Jimenez-Espejo, F.J., et al., 2013. Dynamic behaviour of the East Antarctic Ice Sheet during Pliocene warmth. *Nature Geoscience*, 6(9):765–769. <https://doi.org/10.1038/ngeo1889>
- Cortese, G., Gersonde, R., Hillenbrand, C.-D., and Kuhn, G., 2004. Opal sedimentation shifts in the world ocean over the last 15 Myr. *Earth and Planetary Science Letters*, 224(3–4):509–527. <https://doi.org/10.1016/j.epsl.2004.05.035>
- de Baar, H.J.W., de Jong, J.T.M., Bakker, D.C.E., Löscher, B.M., Veth, C., Bathmann, U., and Smetacek, V., 1995. Importance of iron for plankton blooms and carbon dioxide drawdown in the Southern Ocean. *Nature*, 373(6513):412–415. <https://doi.org/10.1038/373412a0>
- De Deckker, P., Moros, M., Perner, K., and Jansen, E., 2012. Influence of the tropics and southern westerlies on glacial interhemispheric asymmetry. *Nature Geoscience*, 5(4):266–269. <https://doi.org/10.1038/ngeo1431>
- Deacon, G.E.R., 1937. *Discovery Reports (Volume 15): The Hydrology of the Southern Ocean*: Cambridge, United Kingdom (Cambridge University Press).
- Dekens, P.S., Ravelo, A.C., and McCarthy, M.D., 2007. Warm upwelling regions in the Pliocene warm period. *Paleoceanography and Paleoclimatology*, 22(3):PA3211. <https://doi.org/10.1029/2006PA001394>
- Diekmann, B., 2007. Sedimentary patterns in the late Quaternary Southern Ocean. *Deep Sea Research, Part II: Topical Studies in Oceanography*, 54(21–22):2350–2366. <https://doi.org/10.1016/j.dsr.2007.07.025>
- Eagles, G., 2006. Deviations from an ideal thermal subsidence surface in the southern Pacific Ocean. *Terra Antarctica Reports*, 12:109–118.
- Esper, O., and Gersonde, R., 2014. Quaternary surface water temperature estimations: new diatom transfer functions for the Southern Ocean. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 414:1–19. <https://doi.org/10.1016/j.palaeo.2014.08.008>
- Expedition 318 Scientists, 2010. *Expedition 318 Preliminary Report: Wilkes Land Glacial History*. Integrated Ocean Drilling Program. <https://doi.org/10.2204/iodp.pr.318.2010>
- Ferrari, R., Provost, C., Park, Y.-H., Sennéchal, N., Koenig, Z., Sekma, H., Garric, G., and Bourdallé-Badie, R., 2014. Heat fluxes across the Antarctic Circumpolar Current in Drake Passage: mean flow and eddy contributions. *Journal of Geophysical Research: Oceans*, 119(9):6381–6402. <https://doi.org/10.1002/2014JC010201>
- Florindo, F., Nelson, A.E., and Haywood, A.M., 2008. Introduction to “Antarctic cryosphere and Southern Ocean climate evolution (Cenozoic–Holocene).” *Palaeogeography, Palaeoclimatology, Palaeoecology*, 260(1–2):1–7. <https://doi.org/10.1016/j.palaeo.2007.12.001>
- Frank, M., Gersonde, R., van der Loeff, M.R., Bohrmann, G., Nürnberg, C.C., Kubik, P.W., Suter, M., and Mangini, A., 2000. Similar glacial and interglacial export bioproductivity in the Atlantic sector of the Southern Ocean: multiproxy evidence and implications for glacial atmospheric CO₂. *Paleoceanography and Paleoclimatology*, 15(6):642–658. <https://doi.org/10.1029/2000PA000497>
- Gersonde, R., 2011. The Expedition of the Research Vessel “Polarstern” to the Polar South Pacific Sea in 2009/2010 (ANT-XXVI/2 - BIPOMAC). *Berichte zur Polar und Meeresforschung*, 632. <http://epic.awi.de/29941/>
- Gersonde, R., Abelmann, A., Brathauer, U., Becquey, S., Bianchi, C., Cortese, G., Grobe, H., Kuhn, G., Niebler, H.-S., Segl, M., Sieger, R., Zielinski, U., and Fütterer, D.K., 2003. Last glacial sea surface temperatures and sea-ice extent in the Southern Ocean (Atlantic-Indian sector): a multiproxy approach. *Paleoceanography*, 18(3):1061. <https://doi.org/10.1029/2002PA000809>
- Gersonde, R., Crosta, X., Abelmann, A., and Armand, L., 2005. Sea-surface temperature and sea ice distribution of the Southern Ocean at the EPILOG Last Glacial Maximum—a circum-Antarctic view based on siliceous microfossil records. *Quaternary Science Reviews*, 24(7–9):869–896. <https://doi.org/10.1016/j.quascirev.2004.07.015>

- Gordon, A.L., 1986. Inter-ocean exchange of thermocline water. *Journal of Geophysical Research: Oceans*, 91(C4):5037–5046. <https://doi.org/10.1029/JC091iC04p05037>
- Hain, M.P., Sigman, D.M., and Haug, G.H., 2010. Carbon dioxide effects of Antarctic stratification, North Atlantic Intermediate Water formation, and subantarctic nutrient drawdown during the last ice age: diagnosis and synthesis in a geochemical box model. *Global Biogeochemical Cycles*, 24(4):GB4023. <https://doi.org/10.1029/2010GB003790>
- Hall, I.R., Hemming, S.R., and LeVay, L.J., 2015. *Expedition 361 Scientific Prospectus: South African Climates (Agulhas LGM Density Profile)*. International Ocean Discovery Program. <https://doi.org/10.14379/iodp.sp.361.2015>
- Hall, I.R., McCave, I.N., Shackleton, N.J., Weedon, G.P., and Harris, S.E., 2001. Intensified deep Pacific inflow and ventilation in Pleistocene glacial times. *Nature*, 412(6849):809–812. <https://doi.org/10.1038/35090552>
- Hayes, C.T., Martínez-García, A., Hasenfratz, A.P., Jaccard, S.L., Hodell, D.A., Sigman, D.M., Haug, G.H., and Anderson, R.F., 2014. A stagnation event in the deep South Atlantic during the last interglacial period. *Science*, 346(6216):1514–1517. <https://doi.org/10.1126/science.1256620>
- Hilgen, F.J., Lourens, L.J., and Van Dam, J.A., 2012. The Neogene period. With contributions by A.G. Beu, A.F. Boyes, R.A. Cooper, W. Krijgsman, J.G. Ogg, W.E. Piller, and D.S. Wilson. In Gradstein, F.M., Ogg, J.G., Schmitz, M.D., and Ogg, G.M. (Eds.), *The Geologic Time Scale*: Oxford, United Kingdom (Elsevier), 923–978. <https://doi.org/10.1016/B978-0-444-59425-9.00029-9>
- Hillenbrand, C.-D., and Cortese, G., 2006. Polar stratification: a critical view from the Southern Ocean. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 242(3–4):240–252. <https://doi.org/10.1016/j.palaeo.2006.06.001>
- Ho, S.L., Mollenhauer, G., Lamy, F., Martínez-García, A., Mohtadi, M., Gersonde, R., Hebbeln, D., Nunez-Ricardo, S., Rosell-Melé, A., and Tiedemann, R., 2012. Sea surface temperature variability in the Pacific sector of the Southern Ocean over the past 700 kyr. *Paleoceanography and Paleoclimatology*, 27(4):PA4202. <https://doi.org/10.1029/2012PA002317>
- Iversen, N., and Jorgensen, B.B., 1985. Anaerobic methane oxidation rates at the sulfate-methane transition in marine sediments from Kattegat and Skagerrak (Denmark). *Limnology and Oceanography*, 30(5):944–955. <https://doi.org/10.4319/lo.1985.30.5.0944>
- Jaccard, S.L., Hayes, C.T., Martínez-García, A., Hodell, D.A., Anderson, R.F., Sigman, D.M., and Haug, G.H., 2013. Two modes of change in Southern Ocean productivity over the past million years. *Science*, 339(6126):1419–1423. <https://doi.org/10.1126/science.1227545>
- Kaiser, J., and Lamy, F., 2010. Links between Patagonian Ice Sheet fluctuations and Antarctic dust variability during the last glacial period (MIS 4–2). *Quaternary Science Reviews*, 29(11–12):1464–1471. <https://doi.org/10.1016/j.quascirev.2010.03.005>
- Kanfoush, S.L., Hodell, D.A., Charles, C.D., Janecsek, T.R., and Rack, F.R., 2002. Comparison of ice-rafted debris and physical properties in ODP Site 1094 (South Atlantic) with the Vostok ice core over the last four climate cycles. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 182(3–4):329–349. [https://doi.org/10.1016/S0031-0182\(01\)00502-8](https://doi.org/10.1016/S0031-0182(01)00502-8)
- Kawabe, M., and Fujio, S., 2010. Pacific Ocean circulation based on observation. *Journal of Oceanography*, 66(3):389–403. <https://doi.org/10.1007/s10872-010-0034-8>
- Knorr, G., and Lohmann, G., 2003. Southern Ocean origin for the resumption of Atlantic thermohaline circulation during deglaciation. *Nature*, 424(6948):532–536. <https://doi.org/10.1038/nature01855>
- Koenig, Z., Provost, C., Ferrari, R., Sennéchaël, N., and Rio, M.-H., 2014. Volume transport of the Antarctic Circumpolar Current: production and validation of a 20 year long time series obtained from in situ and satellite observations. *Journal of Geophysical Research: Oceans*, 119(8):5407–5433. <https://doi.org/10.1002/2014JC009966>
- Kohfeld, K.E., Graham, R.M., de Boer, A.M., Sime, L.C., Wolff, E.W., Le Quééré, C., and Bopp, L., 2013. Southern Hemisphere westerly wind changes during the Last Glacial Maximum: paleo-data synthesis. *Quaternary Science Reviews*, 68:76–95. <https://doi.org/10.1016/j.quascirev.2013.01.017>
- Kohfeld, K.E., Le Quééré, C., Harrison, S.P., and Anderson, R.F., 2005. Role of marine biology in glacial–interglacial CO₂ cycles. *Science*, 308(5718):74–78. <https://doi.org/10.1126/science.1105375>
- Kuhlbrodt, T., Griesel, A., Montoya, M., Levermann, A., Hofmann, M., and Rahmstorf, S., 2007. On the driving processes of the Atlantic meridional overturning circulation. *Reviews of Geophysics*, 45(2):RG2001. <https://doi.org/10.1029/2004RG000166>
- Kumar, N., Anderson, R.F., Mortlock, R.A., Froelich, P.N., Kubik, P., Dittrich-Hannen, B., and Suter, M., 1995. Increased biological productivity and export production in the glacial Southern Ocean. *Nature*, 378(6558):675–680. <https://doi.org/10.1038/378675a0>
- Lamy, F., 2016. The Expedition PS97 of the Research Vessel POLARSTERN to the Drake Passage in 2016. *Berichte zur Polar und Meeresforschung*, 701. <http://epic.awi.de/41674/>
- Lamy, F., Arz, H.W., Kilian, R., Lange, C.B., Lembke-Jene, L., Wengler, M., Kaiser, J., Baeza-Urrea, O., Hall, I.R., Harada, N., and Tiedemann, R., 2015. Glacial reduction and millennial-scale variations in Drake Passage throughflow. *Proceedings of the National Academy of Sciences of the United States of America*, 112(44):13496–13501. <https://doi.org/10.1073/pnas.1509203112>
- Lamy, F., Gersonde, R., Winckler, G., Esper, O., Jaeschke, A., Kuhn, G., Ullermann, J., Martínez-García, A., Lambert, F., and Kilian, R., 2014. Increased dust deposition in the Pacific Southern Ocean during glacial periods. *Science*, 343(6169):403–407. <https://doi.org/10.1126/science.1245424>
- Lamy, F., Kaiser, J., Arz, H.W., Hebbeln, D., Ninnemann, U., Timm, O., Timmermann, A., and Toggweiler, J.R., 2007. Modulation of the bipolar seesaw in the southeast Pacific during Termination 1. *Earth and Planetary Science Letters*, 259(3–4):400–413. <https://doi.org/10.1016/j.epsl.2007.04.040>
- Lamy, F., Kaiser, J., Ninnemann, U., Hebbeln, D., Arz, H.W., and Stoner, J., 2004. Antarctic timing of surface water changes off Chile and Patagonian Ice Sheet response. *Science*, 304(5679):1959–1962. <https://doi.org/10.1126/science.1097863>
- Lea, D.W., Pak, D.K., Belanger, C.L., Spero, H.J., Hall, M.A., and Shackleton, N.J., 2006. Paleoclimate history of Galápagos surface waters over the last 135,000 yr. *Quaternary Science Reviews*, 25(11–12):1152–1167. <https://doi.org/10.1016/j.quascirev.2005.11.010>
- Liu, Z., Shin, S.-I., Otto-Bliesner, B., Kutzbach, J.E., Brady, E.C., and Lee, D.E., 2002. Tropical cooling at the Last Glacial Maximum and extratropical ocean ventilation. *Geophysical Research Letters*, 29(10):48-1–48-4. <https://doi.org/10.1029/2001GL013938>
- Lorrey, A.M., Vandergoes, M., Almond, P., Renwick, J., Stephens, T., Bostock, H., Mackintosh, A., et al., 2012. Palaeocirculation across New Zealand during the Last Glacial Maximum at ~21 ka. *Quaternary Science Reviews*, 36:189–213. <https://doi.org/10.1016/j.quascirev.2011.09.025>
- Lynch-Stieglitz, J., Ito, T., and Michel, E., 2016. Antarctic density stratification and the strength of the circumpolar current during the Last Glacial Maximum. *Paleoceanography and Paleoclimatology*, 31(5):539–552. <https://doi.org/10.1002/2015PA002915>
- Maksym, T., Stammerjohn, S.E., Ackley, S., and Massom, R., 2012. Antarctic sea ice—a polar opposite? *Oceanography*, 25(3):140–151. <https://doi.org/10.5670/oceanog.2012.88>
- Marshall, J., and Speer, K., 2012. Closure of the meridional overturning circulation through Southern Ocean upwelling. *Nature Geoscience*, 5(3):171–180. <https://doi.org/10.1038/ngeo1391>
- Martin, J.H., 1990. Glacial–interglacial CO₂ change: the iron hypothesis. *Paleoceanography and Paleoclimatology*, 5(1):1–13. <https://doi.org/10.1029/PA005i001p00001>
- Martínez-García, A., Rosell-Melé, A., Geibert, W., Gersonde, R., Masqué, P., Gaspari, V., and Barbante, C., 2009. Links between iron supply, marine productivity, sea surface temperature, and CO₂ over the last 1.1 Ma. *Paleoceanography and Paleoclimatology*, 24(1):PA1207. <https://doi.org/10.1029/2008PA001657>
- Martínez-García, A., Rosell-Melé, A., Jaccard, S.L., Geibert, W., Sigman, D.M., and Haug, G.H., 2011. Southern Ocean dust–climate coupling over the past four million years. *Nature*, 476(7360):312–315. <https://doi.org/10.1038/nature10310>

- Martínez-García, A., Rosell-Melé, A., McClymont, E.L., Gersonde, R., and Haug, G.H., 2010. Subpolar link to the emergence of the modern equatorial Pacific cold tongue. *Science*, 328(5985):1550–1553. <https://doi.org/10.1126/science.1184480>
- Martínez-García, A., Sigman, D.M., Ren, H., Anderson, R.F., Straub, M., Hodell, D.A., Jaccard, S.L., Eglinton, T.I., and Haug, G.H., 2014. Iron fertilization of the Subantarctic Ocean during the Last Ice Age. *Science*, 343(6177):1347–1350. <https://doi.org/10.1126/science.1246848>
- Mazaud, A., Michel, E., Dewilde, F., and Turon, J.L., 2010. Variations of the Antarctic Circumpolar Current intensity during the past 500 ka. *Geochemistry, Geophysics, Geosystems*, 11(8):Q08007. <https://doi.org/10.1029/2010GC003033>
- McCartney, M.S., 1977. Subantarctic mode water. In Angel, M. (Ed.), *A Voyage of Discovery: George Deacon 70th Anniversary Volume*. Deep-Sea Research and Oceanographic Abstracts, 24:103–119.
- McCave, I.N., Crowhurst, S.J., Kuhn, G., Hillenbrand, C.D., and Meredith, M.P., 2014. Minimal change in Antarctic Circumpolar Current flow speed between the last glacial and Holocene. *Nature Geoscience*, 7(2):113–116. <https://doi.org/10.1038/ngeo2037>
- McKay, R., Naish, T., Carter, L., Riesselman, C., Dunbar, R., Sjunneskog, C., Winter, D., et al., 2012. Antarctic and Southern Ocean influences on late Pliocene global cooling. *Proceedings of the National Academy of Sciences of the United States of America*, 109(17):6423–6428. <https://doi.org/10.1073/pnas.1112248109>
- McManus, J.F., Oppo, D.W., and Cullen, J.L., 1999. A 0.5-million-year record of millennial-scale climate variability in the North Atlantic. *Science*, 283(5404):971–975. <https://doi.org/10.1126/science.283.5404.971>
- Meijers, A.J.S., 2014. The Southern Ocean in the Coupled Model Intercomparison Project Phase 5. *Philosophical Transactions of the Royal Society, A: Mathematical, Physical & Engineering Sciences*, 372(2019):20130296. <https://doi.org/10.1098/rsta.2013.0296>
- Meredith, M.P., Woodworth, P.L., Chereskin, T.K., Marshall, D.P., Allison, L.C., Bigg, G.R., Donohue, K., et al., 2011. Sustained monitoring of the Southern Ocean at Drake Passage: past achievements and future priorities. *Reviews of Geophysics*, 49(4):RG4005. <https://doi.org/10.1029/2010RG000348>
- Naafs, B.D.A., Hefter, J., and Stein R., 2013. Millennial-scale ice rafting events and Hudson Strait Heinrich(-like) events during the late Pliocene and Pleistocene: a review. *Quaternary Science Reviews*, 80:1–28. <https://doi.org/10.1016/j.quascirev.2013.08.014>
- Naish, T.R., Powell, R., Levy, R., Wilson, G., Scherer, R., Talarico, F., Krissek, L., et al., 2009. Obliquity-paced Pliocene West Antarctic Ice Sheet oscillations. *Nature*, 458(7236):322–329. <https://doi.org/10.1038/nature07867>
- Orsi, A.H., Whitworth III, T., and Nowlin, W.D., Jr., 1995. On the meridional extent and fronts of the Antarctic Circumpolar Current. *Deep-Sea Research, Part I*, 42(5):641–673. [https://doi.org/10.1016/0967-0637\(95\)00021-W](https://doi.org/10.1016/0967-0637(95)00021-W)
- Pahnke, K., Zahn, R., Elderfield, H., and Schulz, M., 2003. 340,000-year centennial-scale marine record of Southern Hemisphere climatic oscillation. *Science*, 301(5635):948–952. <https://doi.org/10.1126/science.1084451>
- Paolo, F.S., Fricker, H.A., and Padman, L., 2015. Volume loss from Antarctic ice shelves is accelerating. *Science*, 348(6232):327–331. <https://doi.org/10.1126/science.aaa0940>
- Parrenin, F., Masson-Delmotte, V., Köhler, P., Raynaud, D., Paillard, D., Schwander, J., Barbante, C., Landais, A., Wegner, A., and Jouzel, J., 2013. Synchronous change of atmospheric CO₂ and Antarctic temperature during the Last Deglacial Warming. *Science*, 339(6123):1060–1063. <https://doi.org/10.1126/science.1226368>
- Pena, L.D., Cacho, I., Ferretti, P., and Hall, M.A., 2008. El Niño–Southern Oscillation-like variability during glacial terminations and interlatitudinal teleconnections. *Paleoceanography and Paleoclimatology*, 23(3):PA3101. <https://doi.org/10.1029/2008PA001620>
- Pitman, W.C., and Heitzler, J.R., 1966. Magnetic anomalies over the Pacific–Antarctic Ridge. *Science*, 154(3753):1164–1171. <https://doi.org/10.1126/science.154.3753.1164>
- Pollard, D., and DeConto, R.M., 2009. Modelling West Antarctic Ice Sheet growth and collapse through the past five million years. *Nature*, 458(7236):329–332. <https://doi.org/10.1038/nature07809>
- Polonia, A., Torelli, L., Brancolini, G., and Loreto, M.-F., 2007. Tectonic accretion versus erosion along the southern Chile Trench: oblique subduction and margin segmentation. *Tectonics*, 26(3):TC1983. <https://doi.org/10.1029/2006TC001983>
- Reid, J.L., and Lynn, R.J., 1971. On the influence of the Norwegian–Greenland and Weddell Seas upon the bottom waters of the Indian and Pacific Oceans. *Deep Sea Research and Oceanographic Abstracts*, 18(11):1063–1088. [https://doi.org/10.1016/0011-7471\(71\)90094-5](https://doi.org/10.1016/0011-7471(71)90094-5)
- Renault, A., Provost, C., Sennéchal, N., Barré, N., and Kartavtseff, A., 2011. Two full-depth velocity sections in the Drake Passage in 2006—transport estimates. *Deep Sea Research, Part II: Topical Studies in Oceanography*, 58(25–26):2572–2591. <https://doi.org/10.1016/j.dsr2.2011.01.004>
- Reynolds, R.W., Rayner, N.A., Smith, T.M., Stokes, D.C., and Wang, W., 2002. An improved in situ and satellite SST analysis for climate. *Journal of Climate*, 15(13):1609–1625. [https://doi.org/10.1175/1520-0442\(2002\)015<1609:AISAS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1609:AISAS>2.0.CO;2)
- Reynolds, R.W., Smith, T.M., Liu, C., Chelton, D.B., Casey, K.S., and Schlax, M.G., 2007. Daily high-resolution-blended analyses for sea surface temperature. *Journal of Climate*, 20(22):5473–5496. <https://doi.org/10.1175/2007JCLI1824.1>
- Rincon-Martinez, D., Contreras, S., Lamy, F., and Tiedemann, R., 2009. Are glacial (interglacial) in the easternmost Pacific Ocean drier (wetter) during the last 300 000 years? In *Awards Ceremony Speeches and Abstracts of the 19th Annual V.M. Goldschmidt Conference*. *Geochimica et Cosmochimica Acta*, 73(13S):A1102. <https://doi.org/10.1016/j.gca.2009.05.014>
- Robinson, R.S., and Sigman, D.M., 2008. Nitrogen isotopic evidence for a poleward decrease in surface nitrate within the ice age Antarctic. *Quaternary Science Reviews*, 27(9–10):1076–1090. <https://doi.org/10.1016/j.quascirev.2008.02.005>
- Ronge, T.A., Tiedemann, R., Lamy, F., Köhler, P., Alloway, B.V., De Pol-Holz, R., Pahnke, K., Southon, J., and Wacker, L., 2016. Radiocarbon constraints on the extent and evolution of the South Pacific glacial carbon pool. *Nature Communications*, 7:11487. <https://doi.org/10.1038/ncomms11487>
- Sigman, D.M., and Boyle, E.A., 2000. Glacial/interglacial variations in atmospheric carbon dioxide. *Nature*, 407(6806):859–869. <https://doi.org/10.1038/35038000>
- Sigman, D.M., Hain, M.P., and Haug, G.H., 2010. The polar ocean and glacial cycles in atmospheric CO₂ concentration. *Nature*, 466(7302):47–55. <https://doi.org/10.1038/nature09149>
- Sigman, D.M., Jaccard, S.L., and Haug, G.H., 2004. Polar ocean stratification in a cold climate. *Nature*, 428(6978):59–63. <https://doi.org/10.1038/nature02357>
- Sloyan, B.M., and Rintoul, S.R., 2001. Circulation, renewal, and modification of Antarctic Mode and Intermediate Water. *Journal of Physical Oceanography*, 31(4):1005–1030. [https://doi.org/10.1175/1520-0485\(2001\)031<1005:CRAMO>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<1005:CRAMO>2.0.CO;2)
- Sokolov, S., and Rintoul, S.R., 2009. Circumpolar structure and distribution of the Antarctic Circumpolar Current fronts: 1. Mean circumpolar paths. *Journal of Geophysical Research: Oceans*, 114(C11):C11018. <https://doi.org/10.1029/2008JC005108>
- Stammerjohn, S., Massom, R., Rind, D., and Martinson, D., 2012. Regions of rapid sea ice change: an inter-hemispheric seasonal comparison. *Geophysical Research Letters*, 39(6):L06501. <https://doi.org/10.1029/2012GL050874>
- Steig, E.J., Ding, Q., Battisti, D.S., and Jenkins, A., 2012. Tropical forcing of Circumpolar Deep Water inflow and outlet glacier thinning in the Amundsen Sea Embayment, West Antarctica. *Annals of Glaciology*, 53(60):19–28. <https://doi.org/10.3189/2012AoG60A110>
- Studer, A.S., Sigman, D.M., Martínez-García, A., Benz, V., Winckler, G., Kuhn, G., Esper, O., et al., 2015. Antarctic Zone nutrient conditions during the last two glacial cycles. *Paleoceanography and Paleoclimatology*, 30(7):845–862. <https://doi.org/10.1002/2014PA002745>

- Sugden, D.E., McCulloch, R.D., Bory, A.J.-M., and Hein, A.S., 2009. Influence of Patagonian glaciers on Antarctic dust deposition during the last glacial period. *Nature Geoscience*, 2(4):281–285. <https://doi.org/10.1038/ngeo474>
- Talley, L.D., 2013. Closure of the global overturning circulation through the Indian, Pacific, and Southern Oceans: schematics and transports. *Oceanography*, 26(1):80–97. <https://doi.org/10.5670/oceanog.2013.07>
- Teitler, L., Florindo, F., Warnke, D.A., Filippelli, G.M., Kupp, G., and Taylor, B., 2015. Antarctic Ice Sheet response to a long warm interval across marine isotope Stage 31: a cross-latitudinal study of iceberg-rafted debris. *Earth and Planetary Science Letters*, 409:109–119. <https://doi.org/10.1016/j.epsl.2014.10.037>
- Teitler, L., Warnke, D.A., Venz, K.C., Hodell, D.A., Becquey, S., Gersonde, R., and Teitler, W., 2010. Determination of Antarctic Ice Sheet stability over the last ~500 ka through a study of iceberg-rafted debris. *Paleoceanography and Paleoclimatology*, 25(1). <https://doi.org/10.1029/2008PA001691>
- Thompson, D.W.J., Solomon, S., Kushner, P.J., England, M.H., Grise, K.M., and Karoly, D.J., 2011. Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change. *Nature Geoscience*, 4(11):741–749. <https://doi.org/10.1038/ngeo1296>
- Turner, J., Comiso, J.C., Marshall, G.J., Lachlan-Cope, T.A., Bracegirdle, T., Maksym, T., Meredith, M.P., Wang, Z., and Orr, A., 2009. Non-annular atmospheric circulation change induced by stratospheric ozone depletion and its role in the recent increase of Antarctic sea ice extent. *Geophysical Research Letters*, 36(8):L08502. <https://doi.org/10.1029/2009GL037524>
- Ullermann, J., Lamy, F., Ninnemann, U., Lembke-Jene, L., Gersonde, R., and Tiedemann, R., 2016. Pacific-Atlantic Circumpolar Deep Water coupling during the last 500 ka. *Paleoceanography and Paleoclimatology*, 31(6):639–650. <https://doi.org/10.1002/2016PA002932>
- Venz, K.A., and Hodell, D.A., 2002. New evidence for changes in Plio–Pleistocene deep water circulation from Southern Ocean ODP Leg 177 Site 1090. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 182(3–4):197–220. [https://doi.org/10.1016/S0031-0182\(01\)00496-5](https://doi.org/10.1016/S0031-0182(01)00496-5)
- Well, R., and Roether, W., 2003. Neon distribution in South Atlantic and South Pacific waters. *Deep-Sea Research Part I: Oceanographic Research Papers*, 50(6):721–735. [https://doi.org/10.1016/S0967-0637\(03\)00058-X](https://doi.org/10.1016/S0967-0637(03)00058-X)
- Well, R., Roether, W., and Stevens, D.P., 2003. An additional deep-water mass in Drake Passage as revealed by ³He data. *Deep Sea Research, Part I: Oceanographic Research Papers*, 50(9):1079–1098. [https://doi.org/10.1016/S0967-0637\(03\)00050-5](https://doi.org/10.1016/S0967-0637(03)00050-5)
- Yuan, X., 2004. ENSO-related impacts on Antarctic sea ice: a synthesis of phenomenon and mechanisms. *Antarctic Science*, 16(4): 415–425. <https://doi.org/10.1017/S0954102004002238>

Table T1. Expedition 383 hole summary. — = no data.

Hole	Latitude	Longitude	Water depth (m)	Total penetration (m)	Drilled interval (m)	Cored interval (m)	Core recovered (m)	Recovery (%)
U1539A	56°09.0600'S	115°08.0461'W	4071.08	107.6	—	107.6	104.75	97.35
U1539B	56°09.0587'S	115°08.0276'W	4070.45	28.2	—	28.2	21.92	77.73
U1539C	56°09.0711'S	115°08.0285'W	4070.15	268.1	—	268.1	247.50	92.32
U1539D	56°09.0720'S	115°08.0470'W	4070.13	198.0	68.7	129.3	128.18	99.13
Site U1539 totals:			601.9	68.7	533.2	502.35	94.21	
U1540A	55°8.4674'S	114°50.5188'W	3584.60	150.0	—	150.0	155.13	103.42
U1540B	55°8.4656'S	114°50.4985'W	3579.98	150.0	10.0	140.0	140.78	100.56
U1540C	55°8.4474'S	114°50.4994'W	3579.26	6.8	—	6.8	6.77	99.56
U1540D	55°8.4766'S	114°50.5196'W	3577.16	211.0	58.7	152.3	151.02	99.16
U1540E	55°8.4772'S	114°50.5375'W	3577.16	213.0	137.0	76.0	78.87	103.78
Site U1540 totals:			730.8	205.7	525.1	532.57	101.42	
U1541A	54°12.7560'S	125°25.5480'W	3606.32	9.5	—	9.5	9.67	101.79
U1541B	54°12.7553'S	125°25.5431'W	3603.72	138.5	—	138.5	129.27	93.34
U1541C	54°12.7566'S	125°25.5288'W	3602.62	118.1	—	118.1	100.37	84.99
Site U1541 totals:			266.1	0.0	266.1	239.31	89.93	
U1542A	52°42.2880'S	75°35.7922'W	1099.81	169.5	—	169.5	181.74	107.22
U1542B	52°42.2893'S	75°35.7551'W	1101.11	1.4	—	1.4	1.46	104.29
U1542C	52°42.2893'S	75°35.7551'W	1100.21	234.0	9.0	225.0	236.86	105.27
U1542D	52°42.3001'S	75°35.7742'W	1100.71	213.7	11.0	202.7	205.04	101.15
Site U1542 totals:			618.6	20.0	598.6	625.10	104.42	
U1543A	54°35.0631'S	76°40.5900'W	3863.36	339.6	—	339.6	350.62	103.24
U1543B	54°35.0646'S	76°40.5697'W	3865.29	286.5	3.4	283.1	294.97	104.19
Site U1543 totals:			626.1	3.4	622.7	645.59	103.67	
U1544A	55°32.2192'S	71°35.6194'W	2089.86	106.0	3.0	103.0	91.34	88.68
Site U1544 totals:			106.0	3.0	103.0	91.34	88.68	
Expedition 383 totals:			2949.5	300.8	2648.7	2636.26	99.53	

Hole	Total cores (N)	APC cores (N)	HLAPC cores (N)	XCB cores (N)	Date started	Start time UTC (h)	Date finished	End time UTC (h)	Time on hole (hours)	Time on hole (days)
U1539A	12	12	0	0	31 May 2019	1515	2 Jun 2019	0530	38.16	1.59
U1539B	3	3	0	0	2 Jun 2019	0530	3 Jun 2019	1300	31.44	1.31
U1539C	32	26	6	0	3 Jun 2019	1300	5 Jun 2019	1530	50.40	2.10
U1539D	14	14	0	0	5 Jun 2019	1530	7 Jun 2019	0330	36.00	1.50
Site U1539 totals:									156.00	6.50
U1540A	16	16	0	0	7 Jun 2019	1215	9 Jun 2019	1200	47.76	1.99
U1540B	16	16	0	0	9 Jun 2019	1200	10 Jun 2019	1300	24.96	1.04
U1540C	1	1	0	0	10 Jun 2019	1300	10 Jun 2019	1430	1.44	0.06
U1540D	19	18	0	1	10 Jun 2019	1430	11 Jun 2019	2015	29.76	1.24
U1540E	8	8	0	0	11 Jun 2019	2015	12 Jun 2019	2200	25.68	1.07
Site U1540 totals:									129.60	5.40
U1541A	1	1	0	0	17 Jun 2019	1115	18 Jun 2019	0300	15.84	0.66
U1541B	16	15	0	1	18 Jun 2019	0300	19 Jun 2019	0400	24.96	1.04
U1541C	13	13	0	0	19 Jun 2019	0400	20 Jun 2019	0500	24.96	1.04
Site U1541 totals:									65.76	2.74
U1542A	20	19	1	0	3 Jul 2019	2330	5 Jul 2019	0300	27.60	1.15
U1542B	1	1	0	0	5 Jul 2019	1845	5 Jul 2019	2100	2.16	0.09
U1542C	28	28	0	0	5 Jul 2019	2100	7 Jul 2019	0715	34.32	1.43
U1542D	24	24	0	0	7 Jul 2019	0715	8 Jul 2019	0900	25.68	1.07
Site U1542 totals:									89.76	3.74
U1543A	36	36	0	0	8 Jul 2019	2015	11 Jul 2019	2130	73.20	3.05
U1543B	31	31	0	0	11 Jul 2019	2130	14 Jul 2019	0330	54.00	2.25
Site U1543 totals:									127.20	5.30
U1544A	19	12	6	1	14 Jul 2019	2045	16 Jul 2019	1300	40.32	1.68
Site U1544 totals:									40.32	1.68
Expedition 383 totals:									608.64	25.36

Figure F1. Antarctic Circumpolar Current (from Marshall and Speer, 2012) and Expedition 383 sites. Sites U1539 and U1541–1544 are primary sites; Site U1540 is an alternate site. Red circles = planned Expedition 383 sites. Orange = Subantarctic Front (SAF) and Polar Front (PF); line thickness represents variability in latitudinal position of the corresponding front. Green arrows = observed speed and direction of surface ocean currents measured by drifters floating at 15 m water depth. Red-shaded area = estimated glacial dust supply based on reconstructions from Lamy et al. (2014).




Figure F2. Examples of surface buoy trajectories (circles = position every 30 days) indicating northeast flow of northern Antarctic Circumpolar Current (ACC) water after crossing the East Pacific Rise. Also shown is the bifurcation of surface water close to the Chile coast (at ~45°S) with northward flowing water in the Humboldt Current System (HCS) and strongly accelerated southward flow in the Cape Horn Current (CHC) toward the Drake Passage. West–east drifting buoys follow the South Pacific Current (SPC). Modified from Chaigneau and Pizarro (2005) and Lamy et al., 2015.




Figure F3. Basement ages of the South Pacific oceanic crust (from Eagles, 2006; contours indicate age in Ma).




Figure F4. Tectonic setting of the Expedition 383 Chile margin sites and Seismic Line IT95-171 across the southern Chile continental margin. Site U1544 and alternate Proposed Site CHI-6A are located in the fore-arc basin at the upper margin. Figures modified from Polonia et al. (2007).




Figure F5. Central and eastern South Pacific paleoceanography and Drake Passage throughflow (modified from Lamy et al., 2015). The pronounced glacial cooling in the eastern Subantarctic is consistent with a northward extension of Antarctic cold-water influence. Reduced Cape Horn Current (CHC) and more sluggish glacial northern Drake Passage transport decreases the export of Pacific ACC water into the South Atlantic (cold-water route). Reduced Southern Westerly Winds (SWW) core and extended sea ice diminish the wind forcing on the ACC and thus the Drake Passage transport. Stronger winds in the northern SWW enhance the South Pacific Gyre and the export of northern ACC water into the HCS. SAF = Subantarctic Front, APF = Antarctic Polar Front, SACCf = Southern ACC Front, HCS = Humboldt Current System, SPC = South Pacific Current. A. Reconstructed Drake Passage throughflow during the Last Glacial Maximum (LGM) compared to modern setting. B. Schematic view of the Drake Passage region with major surface and intermediate water circulation. C. LGM sea-surface temperature anomalies.




Figure F6. Expedition 383 sites in the Pacific Antarctic Circumpolar Current (ACC) (red circles: site survey cores are in parentheses) and cores previously collected during various expeditions in the Southeast Pacific (white circles; Ocean Drilling Program [ODP] Leg 202 Sites 1232–1240 and Leg 178 Sites 1095–1102) in context of modern oceanography. WSI = winter sea ice, SAF = Subantarctic Front (modern locations after Orsi et al., 1995, and Reynolds et al., 2002, 2007). Dotted lines = inferred Last Glacial Maximum positions. SPC = South Pacific Current, HCS = Humboldt Current System, CHC = Cape Horn Current. Inset figures show vertical water mass structure along two transects in the central and eastern South Pacific (oxygen content: AABW = Antarctic Bottom Water, AAIW = Antarctic Intermediate Water, CDW = Circumpolar Deep Water, PDW = Pacific Deep Water). The geographic locations and water depths of the sites allow us to (1) compare the dynamics of the ACC laterally between the central South Pacific in the vicinity of the East Pacific Rise (EPR) (bathymetric constraints) and the eastern South Pacific before entering the Drake Passage and (2) investigate the vertical structure of the ACC. The sites sample the major water masses of the ACC in the Pacific sector of the Southern Ocean from AAIW (Site U1542), across the CDW/PDW (Sites U1539 and U1541), and potentially down to AABW (U1543), at least for glacial times (Ferrari et al., 2014).




Figure F7. Last Glacial Maximum (LGM) winter SSTs and sea ice extent in the Pacific Southern Ocean (from Benz et al., 2016). Color scale of core symbols represents the same color code as the World Ocean Atlas 2009 winter sea-surface temperatures (WOA09 WSST). Eastern Last Glacial Maximum winter sea ice (E-LGM WSI) estimates include the maximum winter sea ice extent (>15% September concentration) and the average sea ice concentration (40% concentration). Modern winter sea ice edges after Reynolds et al. (2007). GSSTF = Glacial Southern Subtropical Front.




Figure F8. Plio/Pleistocene sea-surface temperature (SST) changes in the subpolar North Pacific (Ocean Drilling Program [ODP] Leg 145 Site 882) and Atlantic Southern Ocean (ODP Leg 177 Site 1090) compared to records from the western (ODP Leg 130 Site 806) and eastern (ODP Leg 138 Site 847) equatorial Pacific. Strong enhancement of meridional SST gradients and shift in orbital frequencies at Site 1090 coincide with the strengthening of the “cold tongue” between 1.2 and 1.8 Ma (from Martínez-García et al., 2010).




Figure F9. Schematic representation of the Antarctic Circumpolar Current (ACC) overturning circulation in the Pacific sector of the Southern Ocean (modified from Ronge et al., 2016). Glacial pattern (left): northernmost extent of sea ice and Southern Westerly Winds (SWW). Increased Antarctic Bottom Water (AABW) salinity by brine rejection and reduced Southern Ocean upwelling favors stratification. Increased dust input promotes primary production and drawdown of CO₂. Modern and deglacial pattern (right): upwelling induced by southward shift of Antarctic sea ice and SWW. The erosion of the deep-water carbon pool releases ¹⁴C-depleted CO₂ toward the atmosphere. Following air–sea gas exchange, the outgassing signal is incorporated into newly formed Antarctic Intermediate Water (AAIW; light blue shading). Blue shading: poorly ventilated old and CO₂-rich waters. Darkest shading 2500–3600 m: water level influenced by hydrothermal CO₂. Green arrows = intermediate water, orange arrows = deep water, light blue areas = sea ice, circular arrows = diffusional and diapycnal mixing.




Figure F10. Left: contrasting paleoproductivity pattern in the Antarctic Zone (AZ; ODP Leg 177 Site 1094) versus the Subantarctic Zone (SAZ; ODP Leg 177 Site 1090) over the past 1 My (from Jaccard et al., 2013). Red and blue shading highlights intervals where AZ and SAZ processes, respectively, are dominantly controlling the partitioning of CO₂ between the ocean interior and the atmosphere. During glacial inception, the first half of the pCO₂ reduction is essentially accomplished by decreasing vertical mixing and upwelling in the AZ (red shading). The second portion of the pCO₂ reduction (blue shading), initiated around 225 ppmv, is achieved by enhancing carbon sequestration resulting from increased iron fertilization in the SAZ, thereby leading the climate system to reach full glacial conditions. Right: records of Subantarctic dust-borne iron flux, phytoplankton productivity, surface nitrate consumption, and atmospheric CO₂ over the last glacial cycle (from Martínez-García et al., 2014). The gray-shaded vertical bars highlight maxima in dust flux that correspond to minima in atmospheric CO₂.




Figure F11. Changes in lithogenic mass accumulation rates (MAR_{Litho}) in the Pacific Southern Ocean (from Lamy et al., 2014). Open diamonds (A–C) = ^{230}Th -normalized MAR_{Litho} . A. Dust MAR in the EDC ice core. B. MAR_{Litho} Core PS75/76-2. C. MAR_{Litho} Core PS75/59-2. D. MAR_{Litho} ODP Leg 177 Site 1090. E. ^{230}Th -normalized MAR_{Litho} values from Cores PS75/59-2, PS75/76-2, E11-2, and E33-2. F. ^{230}Th -normalized C_{29} and C_{31} n -alkane MAR from Core PS75/59-2. G. ^{230}Th -normalized MAR_{Litho} Core PS2489-2.




Figure F12. Oceanographic and bathymetric setting, Site U1539. A. Marine geological features and oceanic fronts (after Orsi et al., 1995) (EPR = East Pacific Rise, FZ = Fracture Zone, SAF = Subantarctic Front, PF = Polar Front). B. Detailed bathymetry with seismic lines and shot points.




Figure F13. (A) Multichannel seismic and (B) Parasound profiles, Site U1539.




Figure F14. Modern salinity and oxygen distribution in the central South Pacific used to visualize major water masses. NPDW = North Pacific Deep Water, LCDW = Lower Circumpolar Deep Water, AABW = Antarctic Bottom Water.




Figure F15. Hole summaries, Site U1539. GRA = gamma ray attenuation, MSP = point magnetic susceptibility, RGB = red green blue, NGR = natural gamma radiation. (Continued on next three pages.)




Figure F15 (continued). (Continued on next page.)




Figure F15 (continued). (Continued on next page.)




Figure F15 (continued).




Figure F16. Summary of primary lithostratigraphic variations, Site U1539. The unit boundary at ~125 m CCSF-A separates high diatom mat (Lithofacies 1b) abundance of the early part of the record from the more frequent and thicker nanofossil ooze facies (Lithofacies 3 and 4) in the latter part of the record. Relative ages of the units are based on preliminary shipboard age model. RGB = red green blue.




Figure F17. Relationships between bulk sedimentary carbonate content and (A) RGB blue and (B) color reflectance L*, Holes U1539A (red) and U1539C (gray). Bracketed symbols = data points from the Holocene (Hole U1539A) that were excluded from the regression analysis.




Figure F18. Characteristic variations in major lithology with changes in natural gamma ray (NGR), gamma ray attenuation (GRA) bulk density, magnetic susceptibility (MS), sediment color reflectance (b^* and L^*), and the RGB blue value, Hole U1539A. Lithologies are determined by visual core description, smear slide analyses, and X-ray imaging. Gray bars show intervals of light-colored nannofossil ooze.




Figure F19. Preliminary age model based on biostratigraphic markers, Site U1539.




Figure F20. Oceanographic and bathymetric setting, Site U1540. A. Marine geological features and oceanic fronts (after Orsi et al., 1995). EPR = East Pacific Rise, FZ = Fracture Zone, SAF = Subantarctic Front, PF = Polar Front. B. Detailed bathymetry with seismic lines and shot points.




Figure F21. Parasound profile, Site U1540. TWT = two-way traveltme, SP = shot point.




Figure F22. Hole summaries, Site U1540. Geomagnetic polarity timescale ages along the depth scale are placed at the midpoint of observed reversals. The placement of Pliocene and early Pleistocene stage boundaries in Holes U1540D and U1540E are estimated using the preliminary Site U1540 shipboard age model. GRA = gamma ray attenuation, MSP = point magnetic susceptibility, RGB = red green blue, NGR = natural gamma radiation. (Continued on next four pages.)




Figure F22 (continued). (Continued on next page.)




Figure F22 (continued). (Continued on next page.)




Figure F22 (continued). (Continued on next page.)




Figure F22 (continued).




Figure F23. Summary of primary lithostratigraphic variations, Site U1540. Lithostratigraphic units were defined based on the distribution, occurrence, and composition of major lithofacies. Relative age of the lithostratigraphic units are based on the preliminary shipboard age model. Lithology: green = diatom ooze (Lithofacies 2), blue = calcareous ooze, light brown = clay-bearing biogenic ooze, dark brown = biogenic-bearing clay).




Figure F24. Relationships between bulk sedimentary carbonate content and (A) RGB blue and (B) color reflectance L*, Holes U1540A (red), U1540B (gray), and U1540D (orange). Data points from the Holocene (Hole U1540B) were excluded from the regression analysis.




Figure F25. A–D. Characteristic variations in major lithology and physical properties, Holes U1540A and U1540D. Lithologies were determined by visual core description, smear slide analyses, and X-ray imaging. Gray bars = nannofossil ooze, white bars = diatom ooze, orange bars = presence (C) or dominance (D) of clay. NGR = natural gamma radiation, MS = magnetic susceptibility, GRA = gamma ray attenuation bulk density, RGB = red green blue.




Figure F26. Preliminary age model based on biostratigraphic and paleomagnetic markers, Site U1540. FO-A: first occurrence in Hole U1540A, FO-D: first occurrence in Hole U1540D, LO-A: last occurrence in Hole U1540A, LO-D: last occurrence in Hole U1540D.




Figure F27. Oceanographic and bathymetric setting, Site U1540. A. Marine geological features and oceanic fronts (after Orsi et al., 1995). EPR = East Pacific Rise, FZ = Fracture Zone, SAF = Subantarctic Front, PF = Polar Front. B. Detailed bathymetry with seismic lines and shot points.




Figure F28. (A) Multichannel seismic and (B) Parasound profiles, Site U1541. TWT = two-way traveltme, CDP = common depth point.




Figure F29. Hole summaries, Site U1541. Geomagnetic polarity timescale ages along the depth scale are placed at the midpoint of observed reversals in Holes U1541B and U1541C. The placement of Pliocene and early Pleistocene stage boundaries in Holes U1541B and U1541C are estimated using the preliminary Site U1541 shipboard age model. GRA = gamma ray attenuation, MSP = point magnetic susceptibility, RGB = red green blue, NGR = natural gamma radiation. (Continued on next two pages.)




Figure F29 (continued). (Continued on next page.)




Figure F29 (continued).




Figure F30. Summary of primary lithostratigraphic variations, Site U1541. Lithostratigraphic units were defined based on the distribution, occurrence, and composition of major lithofacies. Relative age of the units is based on the preliminary shipboard age model for Site U1541. Ages on depth scale indicate magnetic reversals. Lithologies: light green = diatom ooze, light blue = diatom-rich/bearing calcareous ooze, dark blue = nannofossil ooze, light brown = clay-bearing to clayey biogenic ooze. MSP = point magnetic susceptibility, RGB = red green blue.




Figure F31. Age-depth relationship based on correlation to the geomagnetic polarity timescale (Cande and Kent, 1995), Holes U1541B and U1541C.




Figure F32. Relationships between bulk sedimentary carbonate content and (A) RGB blue and (B) color reflectance L*, Holes U1541A and U1541B.




Figure F33. A–E. Characteristic variations in major lithology and physical properties, Hole U1541B. Gray bars (A, B) = diatom ooze, light gray bars (C, D) = diatom bearing, orange bars = clay bearing. GRA = gamma ray attenuation bulk density, NGR = natural gamma radiation, MS = magnetic susceptibility, RGB = red green blue.




Figure F34. Preliminary age model based on biostratigraphic and paleomagnetic markers, Site U1541.




Figure F35. Oceanographic and bathymetric setting, Site U1542. A. Marine geological features. Yellow lines = seismic lines available in the region. B. Detailed bathymetry with seismic lines and shot points.




Figure F36. (A) Multichannel seismic and (B) Parasound profiles, Site U1542. TWT = two-way traveltime, CDP = common depth point.




Figure F37. Schematic view of the southern Chile margin and the Drake Passage region with approximate major surface and intermediate water circulation and Expedition 383 and ODP Leg 202 Site 1233 site locations. ACC = Antarctic Circumpolar Current, SPC = South Pacific Current, HCS = Humboldt Current System, CHC = Cape Horn Current, AAIW = Antarctic Intermediate Water, SAF = Subantarctic Front, WSI = winter sea ice.




Figure F38. Modern oxygen distribution in the eastern South Pacific used to visualize major water masses. PDW = Pacific Deep Water, CDW = Circumpolar Deep Water, AABW = Antarctic Bottom Water, AAIW = Antarctic Intermediate Water.




Figure F39. Hole summaries, Site U1542. GRA = gamma ray attenuation, MSP = point magnetic susceptibility, RGB = red green blue, NGR = natural gamma radiation. (Continued on next 3 pages.)




Figure F39 (continued). (Continued on next page.)




Figure F39 (continued). (Continued on next page.)




Figure F39 (continued).




Figure F40. Summary of primary lithostratigraphic variations, Site U1542. Lithologies: brown = silty clay to clayey silt, yellow = sand, gray = carbonate and/or sand-bearing clayey silt, blue = nannofossil ooze, pale yellow = silt-bearing nannofossil ooze. MSP = point magnetic susceptibility, RGB = red green blue.




Figure F41. Characteristic variations in major lithology and physical properties, Hole U1542C. GRA = gamma ray attenuation bulk density, MSP = point magnetic susceptibility, NGR = natural gamma radiation, RGB = red green blue.




Figure F42. Preliminary age model based on biostratigraphic and paleomagnetic markers, Site U1542.




Figure F43. Oceanographic and bathymetric setting, Site U1543. A. Marine geological features. Yellow lines = seismic lines available in the region. B. Detailed bathymetry with seismic lines and shot points.




Figure F44. (A) Multichannel seismic and (B) Parasound profiles, Site U1542. TWT = two-way traveltime, CDP = common depth point.





Figure F45. Hole summaries, Site U1543A. GRA = gamma ray attenuation bulk density, MSP = point magnetic susceptibility, RGB = red blue green, NGR = natural gamma radiation. (Continued on next page.)




Figure 45 (continued).




Figure F46. Summary of primary lithostratigraphic variations, Site U1543. Lithologies: tan = nannofossil ooze, blue = clay-, silt-, and/or diatom-bearing/rich calcareous ooze, orange = siliciclastic and/or carbonate-bearing/rich diatom ooze, white = diatom-bearing clay, gray = clayey silt to silty clay. MSP = point magnetic susceptibility, RGB = red green blue.




Figure F47. Corrected inclination after 15 mT peak AF demagnetization, polarity interpretation, and correlation to the geomagnetic polarity timescale (GPTS; Hilgen et al., 2012), Hole U1543A.




Figure F48. Relationships between bulk sedimentary carbonate content and (A) RGB blue and (B) color reflectance L*, Hole U1543A.




Figure F49. Characteristic variations in major lithology and physical properties, Hole U1543A. Pale orange bars = nannofossil/calcareous ooze (Lithofacies 4 and 10), green bars = diatom ooze (Lithofacies 6), white bars = siliciclastic sediments (Lithofacies 7 and 8). GRA = gamma ray attenuation bulk density, MS = magnetic susceptibility, NGR = natural gamma radiation, RGB = red green blue.




Figure F50. Preliminary age model based on biostratigraphic and paleomagnetic markers, Site U1543.




Figure F51. Oceanographic and bathymetric setting, Site U1544. A. Marine geological features. Yellow lines = seismic lines available in the region. B. Detailed bathymetry with seismic lines and shot points.




Figure F52. (A) Multichannel seismic and (B) Parasound profiles, Site U1544.




Figure F53. Hole summary, Site U1544A. GRA = gamma ray attenuation bulk density, MSP = point magnetic susceptibility, RGB = red green blue, NGR = natural gamma radiation.




Figure F54. Summary of primary lithostratigraphic variations, Site U1544. Lithologies: gray = silty clay, orange = sand, tan = clay-, silt-, and/or diatom-bearing/rich nannofossil ooze, white = biosilica-bearing silty clay.




Figure F55. Characteristic variations in major lithology and physical properties, Hole U1544A. Orange bars = sand (Lithofacies 11), pale orange bars = nannofossil ooze (Lithofacies 4 and 10), white bars = diatom ooze (Lithofacies 6) and siliciclastic sediments (Lithofacies 7 and 8).

