



Table T8. Calcareous nannofossil range chart, Hole C0001E. (See table notes.)

Epoch	Zone	Core, section, interval (cm)	Depth CSF (m)	Abundance	Preservation	<i>Calcidiscus leptoporus</i>	<i>Calcidiscus macintyreii</i>	<i>Ceratolithus cristatus</i>	<i>Coccolithus miopelagicus</i>	<i>Coccolithus pelagicus</i> (without bar)	<i>Discoaster</i> spp.	<i>Emiliana huxleyi</i>	<i>Florispheera profunda</i>	<i>Gephyrocapsa</i> spp. large (>5.5 µm)	<i>Gephyrocapsa</i> spp. medium (>3.5–4 µm)	<i>Gephyrocapsa</i> spp. medium II (>4–5.5 µm)	<i>Gephyrocapsa</i> spp. small (<3.5 µm)	<i>Helicosphaera carteri</i>	<i>Helicosphaera</i> sp.	<i>Helicosphaera sellii</i>	<i>Helicosphaera wallichii</i>	<i>Micula</i> sp.	<i>Oolithothus fragilis</i>	<i>Pontosphaera japonica</i>	<i>Pontosphaera multipora</i>	<i>Pseudoemiliania lacunosa</i>	<i>Pseudoemiliania ovata</i>	<i>Reticulofenestra asanoi</i>	<i>Reticulofenestra minuta</i>	<i>Reticulofenestra pseudoumbilicus</i> (5–7 µm)	<i>Reticulofenestra</i> sp.	<i>Rhabdosphaera clavigera</i>	<i>Scapholithus fossilis</i>	<i>Scyphosphaera</i> spp.	<i>Syracosphaera pulchra</i>	<i>Syracosphaera</i> spp.	<i>Umbellosphaera irregularis</i>	<i>Umbellosphaera tenuis</i>	<i>Umbilicosphaera foliosa</i>	<i>Umbilicosphaera sibogae</i>													
Pleistocene	NN21	315-C0001E-1H-2, 10.0	0.70	A	G	C					A	A	F	C	C	C	F				F	R																						F	F	C	F						
		1H-3, 38.0	1.83	A	G	F		F		F	C	A	A	F	C	C	A	C					C																														
	NN20	1H-CC, 20–25	4.11	A	G	C							A	A	D	A	D	C					C																														
	NN19		2H-CC, 48.5	13.62	A	G			R			R		C	C	A	A	F									F	F																					R	F	R	F	
			3H-CC, 23–28	14.28	A	G								A	C	A	A	R																																			
			4H-CC, 23.5–28.5	32.59	A	M	C							A	C	C	C	D	F									C	F																								
			5H-CC, 44.5	42.12	A	M	F			R				C				D										R	C																								
			6H-CC, 32.5–37.5	51.60	A	G	F							A		C		D	F				F					C	C																								
			7H-CC, 29.5–34.5	61.10	A	G	C							A		A		A										C	A	C																							
			8H-CC, 45.5–50.5	70.63	A	M	C	R				R		A				A										C	A	C	C																						
			9H-CC, 29–34	80.10	A	M	C							C	C	C	C	D										C	A	C	C																						
			10H-CC, 23.5–28.5	89.60	A	M	C							A	A			A											C	C																							
			11H-CC, 26.5–31.5	99.10	R	P								A	F	F		A											F	R																							
	12H-CC, 21–26	108.60	C	G	C					R		C				F	D	R		R							F	R																									
	13H-CC, 25–30	118.10	C	M	C				F			A		C		C	D	F		R							C	C																									

Notes: CSF = core depth below seafloor. Abundance: D = dominant, A = abundant, C = common, F = few, R = rare. Preservation: G = good, M = medium, P = poor.