

Table T13. Planktonic foraminifer range chart, Hole C0001F. (See table notes.)

Epoch	Zone	Core, section, interval (cm)	Depth CSF (m)	Tubularia tubularioides															
				Turcomiliidae				Turcomiliidae				Turcomiliidae				Turcomiliidae			
Pleistocene	N22	315-C0001F-1H-CC, 23.5–28.5	117.50	C	G			A	A	C		+ +	A			+ +			
		2H-CC, 22–27	127.00	A	G			A	A	R		+ +	A			+ +			
		3H-CC, 51–56	131.97	C	M			R	C	C		+ +	A			+ +			
		4H-CC, 12.5–17.5	145.95	C	G			C	C	C		+ +	A			+ +			
		5H-CC, 0–5	155.48	C	M			C	C	+		+ +	A			+ +			
	N21	6H-CC, 21–26	161.50	C	G			R	C	+		+ +	C			+ +			
		7H-CC, 71–76	170.98	C	M			C	C	+		+ +	A			+ +			
		8H-CC, 45–50	179.75	C	M			A	A	R		+ +	A			+ +			
		9H-CC, 27.5–32.5	189.24	R	M			R	R	A		+ +	R			+ +			
		10H-CC, 37–42	195.79	R	P			C	+	+		+ +	A			+ +			
Pliocene	N18	11H-CC, 13–18	200.76	R	P	C	R	A	+ +	+ +	+ +	R	+ +	A		+ +			
		12H-CC, 18–23	202.34	R	P	+ +	+ +		+ +	+ +	+ +	R	+ +			+ +			
		13H-3, 29–33	203.61	R	M	+ +	+ +		+ +	+ +	+ +	R	+ +			+ +			
		13H-4, 44–48	204.16	C	M	A	+	+	+	R	+ +	+ +	R	+ +		A	+ +		
		13H-CC, 20–25	207.16	R	P	+ +	+ +		+ +	+ +	+ +	R	+ +			+ +			
		14H-1, 125–129	208.45	C	M	+ +	C	+	C	+ +	+ +	R	+ +			C	+ +		
		14H-2, 28–32	208.86	C	P	+ +	C	+		+ +	+ +	R	+ +			A	+ +		
		14H-3, 53–57	210.52	R	P	+ +			+ +	+ +	+ +	R	+ +			A	+ +		
		14H-5, 112–116	212.49	C	M	+ +	+ +	R	C	C	+ +	R	+ +			A	+ +		
		14H-CC, 0–5	213.27	C	M	R	+	R	+ +	R	+ +	R	+ +			A	+ +		
Pliocene–? Miocene	?N17b	15H-CC, 29–34	218.89	B	B										R	+ +	A	+ +	R
		16H-CC, 19.5–24.5	219.64	B	B												A	+ +	
		17H-CC, 13–18	220.29	C	G	C	+		C	+	+	A	C	+					
		18H-CC, 0–1	226.17	R	P														
		19H-CC, 33–38	229.75	B	B														

Notes: CSF = core depth below seafloor. Abundance: A = abundant, C = common, R = rare, + = present, B = barren. Preservation: G = good, M = medium, P = poor.

