




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo



Core Photo



Core Photo




Core Photo


Hole U1353B Core 5H, Interval 27.7-35.58 m (core depth below seafloor)

SAND, MUDDY SAND


The core is characterized greenish gray shelly muddy v.f. sand in the uppermost part, and dark greenish gray, well sorted v.f. micaceous sand. The shelly muddy sand is presumed to be cave-in material. The well-sorted sand is generally homogeneous, probably due to liquefaction during drilling. No distinct burrows were observed; ichnofabric index is 1.




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo



Core Photo



Core Photo




Core Photo

Hole U1353B Core 31X, Interval 161.4-161.62 m (core depth below seafloor)

MUDY SAND


Only the core catcher was available for this core. Well-cemented muddy sand clasts and muddy v.f. flow-in sand are present. Well-cemented muddy v.f. - f. sand clasts contain well-oriented bivalve shells (maximum 2 cm long).




U1353B-32X NO RECOVERY




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo



Core Photo



Core Photo



Core Photo


Hole U1353B Core 54H, Interval 245.8-248.36 m (core depth below seafloor)

SHELL HASH, SANDY MUD


The core is dominated by a shell hash. It is made of bivalves mostly < 1 cm in size. A very dark gray v.f. sandy mud is a secondary lithology. It contains common scattered shells, though locally forming intervals a few centimeters thick. The shell hash in the lower part of the core is likely to be flow-in material. The ichnofabric index is 1.



Core Photo



Core Photo



Core Photo


Hole U1353B Core 57H, Interval 251.6-253.55 m (core depth below seafloor)

MUDGY SAND, SHELL HASH


The core is characterized by dark greenish gray micaceous homogeneous muddy sand and shell hash. The muddy sand contains moderately sorted, v f - f (mostly v f) micaceous sand. Rare to abundant shell fragments are scattered throughout the muddy sand. Shell hash at the top of the core is likely cave in. In the shell hash are fragments of Tawara, Stiracolpus, barnacles and echinoid spines. No distinct burrows are observed in the shell hash with minor amounts of bioturbation in the muddy sand.




Core Photo



Core Photo



Core Photo




Core Photo


Hole U1353B Core 61X, Interval 257.7-258.12 m (core depth below seafloor)

MUDDY SAND


The core catcher was only recovered for this core. The sediment is characterized by dark greenish gray micaceous homogeneous muddy v.f. sand. Abundant shell fragments, mostly bivalves, are scattered throughout. No distinct burrow is observed (Ichnofabric index 1). The upper portion is soft and soupy due to drilling disturbance.




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo



Core Photo



Core Photo




Core Photo

Hole U1353B Core 71X, Interval 346-347.16 m (core depth below seafloor)


SANDY MUD

The core consists only of dark greenish gray v.f.-f. (mostly v.f.) homogeneous well sorted micaceous sandy mud. Shells are generally broken, rare and scattered throughout. No distinct burrows are observed.

Depth CSF-A (m)	Core length (cm)	Section	Core image	Graphic lithology	Coring disturbance	Sedimentary structure	Lithological accessories	Ichnofabric	Shipboard samples	Age	Magnetic susceptibility (Instrument units)	Natural gamma radiation (cps)	b*	Color
347	100	1	CC				1 2 3 4 5			10.80 12.80 14.80 16.80 48.00 58.00 63.00	2.60 3.60			



Core Photo





Core Photo


Hole U1353B Core 73X, Interval 365.2-365.74 m (core depth below seafloor)

SANDY MUD


The core consists only of dark gray v.f.-f. (mostly v.f.) homogeneous well sorted micaceous sandy mud. Shells are generally broken and are rare and scattered throughout. No distinct burrows are observed (ichnofabric index is 1).



Core Photo



Core Photo




Core Photo

Hole U1353B Core 76X, Interval 394-394.28 m (core depth below seafloor)

MUDY SAND


Only the CC was recovered. The core consists of dark gray v.f.-f. (mostly v.f.) homogeneous well sorted muddy sand. It contains common and scattered shell fragments throughout. The top 6 cm of the core are soupy. The ichnofabric index is 1.



U1353B-77XNO RECOVERY



Core Photo





Core Photo

Hole U1353B Core 79X, Interval 422.8-423.29 m (core depth below seafloor)

SANDY MUD

Only the CC was recovered. The core consists of dark gray v.f.-f. (mostly v.f.) homogeneous well sorted slightly micaceous sandy mud. It contains rare and scattered shell fragments throughout. The ichnofabric index is 1.





Core Photo


Hole U1353B Core 80X, Interval 432.3-432.68 m (core depth below seafloor)

SANDY MUD

Only the CC was recovered. The core consists of dark gray v.f.-f. (mostly v.f.) homogeneous well sorted slightly micaceous sandy mud. It contains rare and scattered shell fragments, though locally forming intervals a few centimeters thick. The ichnofabric index is 1.



Core Photo




Core Photo

Hole U1353B Core 83X, Interval 461.1-461.3 m (core depth below seafloor)

MUDDY SAND, SHELL HASH

Only the CC was recovered. The core consists of dark greenish gray v.f.-f. (mostly v.f.) homogeneous well sorted slightly micaceous muddy sand. It contains rare and scattered shell fragments. The uppermost part of the core is a shell hash, most likely fall-in material. The ichnofabric index is 1.



U1353B-84X NO RECOVERY





Core Photo

Hole U1353B Core 85X, Interval 480.3-480.69 m (core depth below seafloor)

SANDY MUD

Only the CC was recovered. The core consists of dark gray v.f.-f. (mostly v.f.) homogeneous well sorted slightly micaceous sandy mud. It contains rare and scattered shell fragments. The ichnofabric index is 1.




Core Photo

Hole U1353B Core 86X, Interval 489.9-490.26 m (core depth below seafloor)

SANDY MUD

Only the CC was recovered. The core consists of dark gray v.f.-f. (mostly v.f.) homogeneous well sorted slightly micaceous sandy mud. It contains rare and scattered shell fragments. The ichnofabric index is 1.





Core Photo


Hole U1353B Core 87X, Interval 500-500.21 m (core depth below seafloor)

SANDY MUD


Only the CC was recovered. The core consists of dark gray v.f.-f. (mostly v.f.) homogeneous well sorted slightly micaceous sandy mud. It contains common and scattered shell fragments. There are small calcareous concretions (diameter < 2 cm) at the very top of the core. The ichnofabric index is 1.




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo



Core Photo



Core Photo



Core Photo

