

**Table T6 (continued).**

Core, section, interval (cm)	Top depth CSF-A (m)	Interpreted age	NN zone	NZ stage correlation	NZ stage	Age (Ma)	Planktonic foraminifer (%)	Oceanicity	Key species and comments									
									Key species and comments									
98X-CC	604.60	middle Miocene?	NN6–NN4	ND			0	Sheltered inner neritic	ND, no planktonics; middle Miocene nannofossil									

Notes: ND = not determined. HO = highest occurrence, LO = lowest occurrence. Bold = age diagnostic. For New Zealand (NZ) stage abbreviations, see Figure F5 in the “Methods” chapter. This table is also available in [ASCII](#).

**Table T7.** Planktonic foraminifer abundance, preservation, and distribution, Hole U1353A. ([See table notes](#).)

Core, section, interval (cm)	Top depth CSF-A (m)	Foraminiferal preservation	Planktonic foraminifers	Miscellaneous										Comments	
				Zeegebobergia sp. spp.	Turritellina terebelliformis	Tubothalassinids	Obulina unifasciata	Negligible abundance in core	Neogloborotalita inaequilatera	Globocassidina planuloides	Histrioletta siliqua	Globigerinoides rubescens	Globigerinella sphaerica	Globigerinella sphaerica	
317-U1353A-1H-1, 0	0.00	A G 4	X	X	X X	X X	X X	X X	X X	X X	X X	X X	X X	X X	
1H-CC	5.44	A G 4	X X	X	X X	X X	X X	X X	X X	X X	X X	X X	X X	X X	R A D A A
2H-CC	9.33	A G 5	X	X	X X										F A A C
3H-CC	17.95	A G 2	X	X	X	X	X	X	X	X X	X X	X X	X X	X X	
4H-CC	26.72	A G 4	X X	X	X X	X X	X X	X X	X X	X X	X X	X X	X X	X X	
5H-CC	34.08	A G 7	X X	X	X X	X X	X X	X X	X X	X X	X X	X X	X X	X X	
6H-CC	41.37	A G 8	X X	X	X X	X X	X X	X X	X X	X X	X X	X X	X X	X X	
7H-CC	48.33	A G 15	X X	X	X X	X X	X X	X X	X X	X X	X X	X X	X X	X X	
8H-CC	55.97	A G 8	X X	X ?	X X	X X	X X	X X	X X	X X	X X	X X	X X	X X	

Notes: Abundance: D = dominant, A = abundant, C = common, F = few, R = rare, X = present, ? = questionable presence. Preservation: G = good, M = moderate, P = poor. This table is also available in [ASCII](#).