

Table T9. Planktonic foraminiferal abundance, preservation, and distribution, Hole U1354A. (See table notes.)

Core, section, interval (cm)	Top depth CSF-A (m)	Epoch	NZ stage	Planktonic foraminifers												Miscellaneous	Comments			
				Foraminifera bivalves			Planispirina testaceous			Cibicidoides ruber			Cibicidoidella aequilateralis			Orbulina universa				
				P	A	G	P	A	G	G	?	P	A	G	P	A	G	T	T	T
317-U1354A-1H-1, 0	0.00	Holocene	Wq-uWc		A	G	—		X	X	X	X	X	X	X	X	C	D	A	F
1H-CC	3.78				A	G	3	X	X	X	X	X	X	X	X	X	D	A	D	D
2H-CC	12.26				A	G	2	X	X	X	X	X	X	X	X	X	D	D	C	D
3H-CC	13.99				A	G	43	X	X	X	X	X	X	X	X	X	D	D	C	A
4H-CC	14.30				C	G	<1	X									F	F	F	A
5H-CC	14.69				A	G	1										F	F	R	A
6H-CC	24.25				A	G	51	X	X	X	X	X	X	X	X	X	F	F	R	A
7H-CC	31.33				A	G	15	X	X	X	X	X	X	X	X	X	F	F	R	A
8H-CC	38.14				A	G	22	X	X	X	X	X	X	X	X	X	F	F	R	A
9H-CC	46.88				A	G	12	X	X	X	X	X	X	X	X	X	F	F	R	A
10H-CC	53.76				A	G	5	X	X	X	X	X	X	X	X	X	F	F	R	A
11H-CC	60.14				A	G	2	X	X	X	X	X	X	X	X	X	F	F	R	A
12H-CC	64.90				A	G	2	X	X	X	X	X	X	X	X	X	F	F	R	A
13H-CC	73.44				A	G	<1	X	X	X	X	X	X	X	X	X	F	F	R	A
14H-CC	75.41				A	G	6	X	X	X	X	X	X	X	X	X	F	F	R	A
15H-CC	76.20				A	G	1	X	X	X	X	X	X	X	X	X	F	F	R	A
17H-CC	80.16																			
18H-CC	80.33																			
19H-CC	85.38																			

Notes: Abundance: D = dominant, A = abundant, C = common, F = few, R = rare, X = present, ? = questionable presence. Preservation: G = good, M = moderate, P = poor. — = no data. For New Zealand (NZ) stage abbreviations, see Figure F5 in the "Methods" chapter. This table is also available in [ASCII](#).

