

Table T2. Siliceous microfossil abundance and preservation, Hole U1358A. (See table notes.)

Core, section, interval (cm)	Depth (mbsf)		Diatoms												Sponge spicules										
	Top	Bottom	Abundance		Preservation		Preservation - dissolution		Atrypoceraspis cf. acinoceraspis		Actinocyclus cf. actinochilus		Actinocyclus cf. acinoceraspis		Actinocyclus cf. acinoceraspis										
318-U1358A-1R-CC	0.90	1.10	C	P-M	G	X	X	C	X	X	C	C	X	C	R	X	X	F	X	F	F	X	X	R	X
318-U1358B-1R-1, 0	0.00	0.00	A	M	G	X	X	X	X	X	X	F	F	C	X	C	X	X	R	R	X	X	?	X	
1R-CC	0.23	0.42	C	P	G	X	X	X	X	X	X	F	F	C	X	C	X	X	X	X	X	X	X	X	
2R-CC	9.13	9.32	R	P	G	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
3R-2, 24-25	19.03	19.04	X	P	G	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
3R-CC	21.34	21.54	X	P	G	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
4R-1, 92	27.32	27.32	X	P	G	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
4R-CC	28.38	28.62	X	P	G	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	

Notes: * = early form, see Harwood and Maruyama (1992). Data excludes radiolarians. Abundance: A = abundant, C = common, R = rare, X = trace. Preservation: G = good, M= medium, P = poor, ? = uncertainty. See “[Biostratigraphy](#)” in the “Methods” chapter for abundance and preservation definitions.

