

Table T7. Distribution of diatoms, Site U1336. (See table notes.)

Core, section	Preservation	Abundance	<i>Cestodiscus peplum</i>	<i>Cestodiscus pulchellus</i>	<i>Cestodiscus trochus</i>	<i>Coscinodiscus lewisianus</i> var. <i>spiralis</i>	<i>Crucidentacula nicobarica</i>	<i>Ethmodiscus</i> (fragments)	<i>Hemidiscus cuneiformis</i>	<i>Rhaphidiscus marylandicus</i>	<i>Rocella vigilans</i>	<i>Rosiella paleaceae</i>	<i>Synechra jousefana/miocenica</i>	<i>Thalassionema nitzschioides</i> var. <i>parva</i>	<i>Thalassionema robusta</i>	<i>Thalassiothrix yabei</i>
320-U1336B-																
1H-CC	M	F	?	R			R						F			R
2H-CC	P	R					R								?	
3H-CC	P-M	R						R								
4H-CC		B														
5H-CC	P	R		R				R								
6H-CC	P	R			?										?	
7H-CC	P-M	R		R		?				?				R		
8H-CC		B														
9H-CC		B														
10H-CC		B														
11H-CC		B														
12H-CC		B														
13H-CC		B														
14H-CC		B														
15H-CC		B														
16H-CC		B														
17H-CC		B														
18H-CC		B														
19H-CC		B														
20H-CC		B														

Notes: All samples are of middle Miocene age. Preservation: M = moderate, P = poor. Abundance: F = few, R = rare, B = barren, ? = uncertainty.