

**Table T9.** Distribution of planktonic foraminifers, Hole C0011B. (See table notes.)

Core, section, interval (cm)	Sample code	Depth CSF (m)	Abundance	Preservation	Notes
322-C0011B-3R-CC, 12–17	PAL	364.98	R	M	
4R-CC, 0–5	PAL	368.61	R	M	
6R-CC, 7.5–12.5	PAL	393.40	C	G	
7R-CC, 11.5–16.5	PAL	400.83	R	M	
8R-CC, 0–5	PAL	411.17	A	G	
9R-CC, 16–21	PAL	417.69	C	M	
10R-CC, 0–3	PAL	425.13	R	M	
15R-CC, 0–2	PAL	426.79	R	P	
31R-CC, 0–5	PAL	591.05	R	M	
34R-CC, 0–5	PAL	615.17	R	G	
38R-CC, 9.5–14.5	PAL	662.22	R	G	
40R-CC, 0–5	PAL	676.83	R	G	
43R-CC, 8–13	PAL	689.02	R	G	
44R-CC, 0–5	PAL	697.56	R	M	
46R-1, 87–92	PALW	708.89	R	M	
47R-3, 61–67	PALW	713.82	R	M	
48R-CC, 0–5	PAL	726.71	R	M	
49R-CC, 3.5–8.5	PAL	736.48	R	M	
50R-CC, 35–40	PAL	741.44	R	M	
51R-CC, 15.5–20.5	PAL	756.98	R	M	
52R-CC, 15.5–20.5	PAL	766.49	R	M	
53R-CC, 0–5	PAL	770.12	R	M	
56R-CC, 4–9	PAL	846.00	R	M	

Notes: PAL = paleontologic sample, PALW = additional paleontologic sample. Abundance: A = abundant, C = common, R = rare, + = present, B = barren. Preservation: G = good, M = medium, P = poor. Barren samples: 322-C0011B-1R-CC, 35.5–40.5 cm; 2R-CC, 0–5 cm; 5R-CC, 11–16 cm; 11R-CC, 0–3 cm; 12R-CC, 0–5 cm; 13R-CC, 0–5 cm; 14R-CC, 0–5 cm; 16R-CC, 0–5 cm; 17R-CC, 0–5 cm; 19R-CC, 7.5–12.5 cm; 19R-5, 127–131 cm; 20R-CC, 22–27 cm; 21R-CC, 11–16 cm; 22R-CC, 15–20 cm; 23R-CC, 0–5 cm; 23R-1, 49–53 cm; 24R-CC, 23–28 cm; 26R-CC, 7–12 cm; 27R-CC, 0–5 cm; 28R-CC-bottom 5 cm; 30R-CC, 0–5 cm; 33R-CC, 0–7 cm; 35R-CC, 0–5 cm; 37R-CC, 10.5–15.5 cm; 39R-CC, 0–1 cm; 41R-CC, 15.5–20.5 cm; 42R-CC, 0–5 cm; 45R-7, 136–141 cm; 54R-CC, 35.5–40.5 cm; 55R-CC, 12.5–17.5 cm; 57R-CC, 16–21 cm; 58R-CC, 0–5 cm; 59R-CC, 25–30 cm; 60R-CC, 11–12 cm; 61R-CC, 0–5 cm. No samples: 18R-CC; 25R-CC; 32R-CC.