

Table T14. Radiolarian range chart, Holes U1339A, U1339B, U1339C, and U1339D. (See table notes.) (Continued on next page.)

Core, section	Depth (mbsf)		Abundance	Preservation	<i>Actinomma boreale</i>	<i>Amphimelissa setosa</i>	<i>Axoprunum acquilonium</i>	<i>Botryostrobus aquilonaris</i>	<i>Ceratospyrus borealis</i>	<i>Cycladophora cornutoides</i>	<i>Cycladophora davistiana</i>	<i>Dictyophimus hirundo</i>	<i>Larcopyle buetschlii</i>	<i>Lychnocanoma nipponica sakaii</i>	<i>Sphaeropyle langii-robusta</i> group	<i>Spongodiscus</i> sp. (Ling, 1973)	<i>Spongopyle osculosa</i>	<i>Spongotrochus glacialis</i>	<i>Spongurus pylomaticus</i>	<i>Stylatractus universus</i>	<i>Stylochlamydidium venustum</i>	<i>Stylocictya validispina</i>
	Top	Bottom																				
323-U1339A-																						
1H-CC	4.82	4.94	F	G					F		C		F								F	F
2H-CC	14.64	14.74	F	G	F				F		C	F	F								F	F
3H-CC	24.23	24.33	F	G	F				F		F			F							F	F
4H-CC	33.79	33.89	C	G	F			F	C		F			C							F	F
323-U1339B-																						
1H-CC	3.91	4.01	F	G							F						F				F	F
2H-CC	13.65	13.75	F	G					F		C	F	F				F				F	F
3H-CC	23.36	23.36	F	G					F		F	F	F				F				F	F
4H-CC	32.91	33.01	C	G				F	F		F	F	F	C			F				F	F
5H-CC	42.17	42.27	F	G	F			F			F	F	F	C					F		F	F
6H-CC	52.23	52.33	F	G	F			F			C	F	F	F							F	F
7H-CC	61.50	61.60	F	G							F			F								F
8H-CC	70.85	70.95	F	G	F						C	F	F	F								F
9H-CC	80.37	80.47	F	G	F	F					C			F	F	C		F			F	F
10H-CC	89.88	89.98	F	G	F				F		F			F	F	F		F			F	F
11H-CC	99.25	99.35	F	G							F			F								F
12H-CC	108.40	108.48	F	M	F				F					F	F	F						F
13H-CC	108.89	108.99	F	G	F				F		C		F	F	F	F						F
14H-CC	120.47	120.57	F	G							F	F	F	F			F					F
15H-CC	130.02	130.12	F	G			F				C	F	F			F		F				F
16H-CC	139.41	139.51	F	G			F															F
17H-CC	148.57	148.67	F	G	F			C			F		F	F	F	C						F
18H-CC	158.79	158.89	F	G	F	F		F	F		C	F		F	F	F		F			F	F
19H-CC	167.86	167.96	C	M	F	F	F	C			F			F	F	F		F		F		F
20H-CC	177.31	177.41	C	G	C	F		C	F	C	C	C		C	C	F		F		F		F
21H-CC	187.60	187.70	F	G	C			F	F	A	F	F		C	F	F	F	F	F	F	F	F
22H-CC	196.83	196.93	F	G			F	F	F	A	F	F		F	F	F		F		F		F
323-U1339C-																						
1H-CC	9.68	9.78	C	G	F				C		C	F	F		F		F				F	F
2H-CC	19.42	19.52	C	G	F				F		C	F	F	C			F				F	F
3H-CC	29.09	29.19	C	G	F			F	F		C	F	F	F								F
4H-CC	38.86	38.96	F	G	F	F			F		C			F			F					F
5H-CC	48.39	48.39	C	G		F			F		C	F	F	C	F		F					F
6H-CC	57.84	57.94	C	G		F			C		C	F	F	F								F
7H-CC	66.55	66.65	F	G					F		F		F	F				F				F
8H-CC	76.53	76.63	F	G	F	F			F		F	F	F	F	F			F	F		F	F
9H-CC	85.24	85.34	C	M	F				F		F	F	F			C		F				F
10H-CC	95.01	95.11	F	G		F	F	F	F							C						F
11H-CC	104.43	104.53	C	G	F		F	F	F		C					C		F				F
12H-CC	114.36	114.46	F	G	F	F			F		C				F	F	F	F		F		F
13H-CC	123.71	123.81	C	G	C	F			C			F	F		F	F	F	F				F
14H-CC	133.99	134.09	F	G	F			F							F		F					F
15H-CC	142.92	143.02	F	G							F	F				F					F	F
16H-CC	146.52	146.62	C	G	F	F			F		F					C						F
17H-CC	157.02	157.12	C	G	F	F			F	F	A	F		F	F							F
18H-CC	165.92	166.02	F	G	F	F			F		F	F	F									F
19H-CC	176.38	176.48	F	G		F					F											F
20H-CC	185.75	185.85	C	G	F	F			C	F	C	F	C		F	F	F	F	F	F	F	F
21H-CC	195.24	195.34	C	G					F	F	C				F	C						F
323-U1339D-																						
1H-CC	6.52	6.62	F	G	F				F		C											F
2H-CC	16.04	16.14	C	G					C		C	F		F								C
3H-CC	25.59	25.69	C	G					F	F	A	F	F	F								F
4H-CC	33.38	33.48	F	G		F			C				F	F								F
5H-CC	43.24	43.34	F	G					F		C	F		F	F							F



Table T14 (continued).

Core, section	Depth (mbsf)		Abundance	Preservation	<i>Actinomma boreale</i>	<i>Amphimelissa setosa</i>	<i>Axoprunum acqilonium</i>	<i>Botryostrobus aquilonaris</i>	<i>Ceratospyrus borealis</i>	<i>Cycladophora cornutoidea</i>	<i>Cycladophora davisiana</i>	<i>Dictyophimus hirundo</i>	<i>Larcopyle buetschlii</i>	<i>Lychinocanoma nipponica sakaii</i>	<i>Sphaeropyle langii-robusta</i> group	<i>Spongodiscus</i> sp. (Ling, 1973)	<i>Spongopyle osculosa</i>	<i>Spongotrochus glacialis</i>	<i>Spongurus pylomaticus</i>	<i>Stylatractus universus</i>	<i>Stylochlamydidium venustum</i>	<i>Stylocictya validispina</i>
	Top	Bottom																				
6H-CC	52.97	53.07	C	G	F				C	F	A	F	F	F	F		F			F	C	
7H-CC	62.48	62.58	F	G						F	F				F		F			F	F	
8H-CC	71.20	71.30	F	G	F			F		F	C		F	F			F			F	F	
9H-CC	80.15	80.25	C	G	F		F	C		F	C	F	F	F		C				F	F	
10H-CC	90.43	90.53	F	G				F		F						F						
11H-CC	100.38	100.48	F	G	F			F								F						
12H-CC	110.15	110.25	F	G	F			F		F		F				F		F				
13H-CC	119.89	119.99	C	G		F		F		F		F			F	F	F					
14H-CC	128.83	128.93	C	G		F		F		C					F	F						
15H-CC	137.14	137.24	A	G	F	C	C	C		F		F			C	C	F		F	C		
16H-CC	147.16	147.26	C	G	F	F	F	F		F					F	C	F			F		
17H-CC	157.88	157.98	C	G	F	F		F		F	C				F	C				F	F	
18H-CC	167.18	167.28	F	G	F	F		F		F					F	F	F			F		
19H-CC	176.77	176.87	C	G	F	F		F		F	C		F		F		F					
20H-CC	185.99	186.09	F	G	F	F		F		C					F		F					
21H-CC	194.11	194.21	F	G		F		F				F			F	F					F	
22H-CC	200.77	200.83	F	G		F		F		F		F			F						F	

Notes: Abundance: A = abundant, C = common, F = few. Preservation: G = good, M = moderate.