



Core Photo




Core Photo

Hole 330-U1373A-1R Section 2, Top of Section: 0.8 CSF-A (m)


Major lithology: Matrix-supported, brown basalt breccia with few bioclast. Composition of the inter-pebble and cobble spaces consists of a silty-pebbly basalt sediment with a matrix composed of a pale brown, azoic micrite.
Basalt clasts (Subunit 1B):
Type 1 (moderately plagioclase-olivine-phryic basalt, 3% vesicles, light reddish gray)
Type 2 (aphyric basalt, 0-10% vesicles, brownish gray)
Type 3 (aphyric basalt, 15% vesicles, light gray)
Type 4 (highly olivine-phryic basalt, 3% vesicles, reddish gray)
Type 5 (aphyric basalt, 15% vesicles, reddish medium gray to brownish gray)
Type 6 (moderately plagioclase-phryic basalt, 10% vesicles, reddish medium gray)
Type 7 (highly olivine-augite-phryic basalt, 7% vesicles, medium gray)




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo



Core Photo



Core Photo



Core Photo

Hole 330-U1373A-6R Section 2, Top of Section: 29.17 CSF-A (m)

Major lithology: Matrix-supported, dark multicolor basalt breccia. The matrix of the breccias is composed of finer-grained volcaniclastic sediment.

Basalt clasts (Subunit IIID):

- Type 7 (highly olivine-augite-phryic basalt, 7% vesicles, medium gray)
- Type 10 (aphyric basalt, 15% vesicles, pale orange gray)
- Type 11 (aphyric basalt, 10% vesicles, medium gray)
- Type 12 (aphyric basalt, 0.5 % vesicles, brownish red)

Depth (m)
Core length (cm)
Unit
Core image
Graphic lithology

Magnetic susceptibility ($\times 10^{-5}$ SI)
(MS \blacktriangle , MS Point \bullet
- raw;
MS \blacksquare , MS Point \bullet
- filtered)

Reflectance (L⁻¹ = raw;
L⁻¹ = filtered)
(b⁺ \blacktriangle , a⁺ \square = raw;
b⁺ \blacksquare , a⁺ \blacktriangle = filtered)

Grain size

Bioturbation intensity

Ichnofossil/Fossil

Sedimentary structure

Lithologic accessories

Drilling disturbance

Nannofossil zone

P. foraminiferal zone

Age


Shipboard sample

Subunit IIID


29.2
10
20
29.4
30
29.6
40
50
29.8
60
70
80
90
30.0

MADC
TS


Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo




Core Photo



Core Photo



Core Photo



Core Photo

