



Hole 361-U1479A Core 1H, Interval 0.0-1.01 m (CSF-A)

OOZE, NANNOFOSSIL, FORAMINIFERA Core 1 comprises one lithological unit. Unit 1 is light brown (7.5YR 6/4) to light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479B Core 1H, Interval 0.0-6.25 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 1 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. One sandy interval in Section 2 at 137-145 cm. Moderate drilling disturbance in Section 1.




Hole 361-U1479B Core 2H, Interval 6.2-13.75 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 2 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate to severe drilling disturbance in Sections 2, 3 and 5.



Hole 361-U1479B Core 3H, Interval 15.7-24.15 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 3 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Moderate bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.






Hole 361-U1479B Core 4H, Interval 25.2-33.92 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 4 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe to extreme drilling disturbance in Section 1.




Hole 361-U1479B Core 5H, Interval 34.7-44.41 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 5 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.




Hole 361-U1479B Core 6H, Interval 44.2-53.16 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 6 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate to extreme drilling disturbance in Sections 1-3 and 7.




Hole 361-U1479B Core 7H, Interval 53.7-60.56 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 7 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Slight to extreme drilling disturbance in Section 1.




Hole 361-U1479B Core 8H, Interval 63.2-68.09 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 8 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze. A sandy interval in Section 3 at 80-84 cm. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.




Hole 361-U1479B Core 9H, Interval 72.7-81.64 m (CSF-A)

OOZE, NANNOFOSSILS, Core 9 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 1.




Hole 361-U1479B Core 10H, Interval 82.2-92.11 m (CSF-A)

OOZE, NANNOFOSSILS, Core 10 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.



Hole 361-U1479B Core 11H, Interval 91.7-101.51 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 11 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.






Hole 361-U1479B Core 12H, Interval 101.2-111.06 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 12 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 7.




Hole 361-U1479B Core 13H, Interval 110.7-120.55 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 13 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) to light olive gray (5Y 6/2) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate to severe drilling disturbance in Sections 1, 6 and 7.




Hole 361-U1479B Core 14H, Interval 120.2-129.86 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 14 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.




Hole 361-U1479B Core 15H, Interval 129.7-139.62 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 15 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.




Hole 361-U1479B Core 16H, Interval 139.2-149.02 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 16 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera and quartz. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479B Core 17H, Interval 148.7-158.36 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 17 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera and quartz. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 7.




Hole 361-U1479B Core 18H, Interval 158.2-168.23 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 18 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera and quartz. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.



Hole 361-U1479B Core 19H, Interval 167.7-176.97 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 19 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera and quartz. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.






Hole 361-U1479B Core 20H, Interval 177.2-187.15 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 20 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera and quartz. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479B Core 21H, Interval 186.7-196.73 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 21 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera and quartz. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479B Core 22H, Interval 196.2-206.16 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 22 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera and quartz. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479B Core 23H, Interval 205.7-215.65 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 23 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera and quartz. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 1.




Hole 361-U1479B Core 24H, Interval 215.2-225.05 m (CSF-A)

OOZE, NANNOFOSSILS Core 24 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.




Hole 361-U1479B Core 25H, Interval 224.7-234.59 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 25 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.



Hole 361-U1479B Core 26H, Interval 234.2-244.2 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 26 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 1.





Hole 361-U1479B Core 27H, Interval 243.7-253.37 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 27 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate to severe drilling disturbance in Sections 1-2.






Hole 361-U1479B Core 28H, Interval 253.2-263.22 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 28 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479B Core 29H, Interval 262.7-272.13 m (CSF-A)

OOZE, NANNOFOSSILS Core 29 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY) nannofossil ooze. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate to severe drilling disturbance in Sections 1 and 4.




Hole 361-U1479B Core 30H, Interval 272.2-282.25 m (CSF-A)

OOZE, NANNOFOSSILS Core 30 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY) nannofossil ooze. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.




Hole 361-U1479B Core 31H, Interval 281.7-291.68 m (CSF-A)


OOZE, NANNOFOSSILS Core 31 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY) nannofossil ooze. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.



Hole 361-U1479B Core 32H, Interval 291.2-301.03 m (CSF-A)


OOZE, NANNOFOSSILS Core 32 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY) nannofossil ooze. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Slumps or severe drilling disturbance are observed in all Sections of the Core.






Hole 361-U1479C Core 2H, Interval 1.0-10.57 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 2 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Slight drilling disturbance in Section 1.



Hole 361-U1479C Core 3H, Interval 10.5-20.11 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 3 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Slight drilling disturbance in Section 1.






Hole 361-U1479C Core 4H, Interval 20.0-29.72 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 4 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.




Hole 361-U1479C Core 5H, Interval 29.5-39.12 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 5 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Slight drilling disturbance in Section 1.




Hole 361-U1479C Core 6H, Interval 39.0-48.73 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 6 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. One sandy interval in Section 6 at 36-40 cm and one in Section 7 at 22-24 cm. Severe drilling disturbance in Section 1.




Hole 361-U1479C Core 7H, Interval 48.5-58.14 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 7 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.




Hole 361-U1479C Core 8H, Interval 58.0-67.62 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 8 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. One sandy interval in Section 7 at 18-21.5 cm. Extreme drilling disturbance in Section 1.




Hole 361-U1479C Core 9H, Interval 67.5-77.0 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 9 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.




Hole 361-U1479C Core 10H, Interval 77.0-86.89 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 10 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.



Hole 361-U1479C Core 11H, Interval 86.5-96.42 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 11 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.






Hole 361-U1479C Core 12H, Interval 96.0-105.99 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 12 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.




Hole 361-U1479C Core 13H, Interval 105.5-115.51 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 13 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.




Hole 361-U1479C Core 14H, Interval 115.0-124.97 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 14 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) to light olive gray (5Y 6/2) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Sections 1 and 7.




Hole 361-U1479C Core 15H, Interval 124.5-134.47 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 15 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479C Core 16H, Interval 134.0-144.03 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 16 comprises one lithological unit. Unit 1 is light greenish gray (GLE Y 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.




Hole 361-U1479C Core 17H, Interval 143.5-153.53 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 17 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.



Hole 361-U1479C Core 18H, Interval 153.0-161.75 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 18 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.









Hole 361-U1479C Core 20H, Interval 172.0-181.88 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 20 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 7.




Hole 361-U1479C Core 21H, Interval 181.5-191.51 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 21 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479C Core 22H, Interval 191.0-200.96 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 22 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Sections 6 and 7.




Hole 361-U1479C Core 23H, Interval 200.5-210.4 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 23 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1.




Hole 361-U1479C Core 24H, Interval 210.0-219.43 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 24 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 1.




Hole 361-U1479C Core 25H, Interval 219.5-229.07 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 25 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 7.




Hole 361-U1479C Core 26H, Interval 229.0-239.01 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 26 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Slight drilling disturbance in Section 1.



Hole 361-U1479C Core 27H, Interval 238.5-248.27 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 27 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Moderate to strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 1.






Hole 361-U1479C Core 28H, Interval 248.0-257.9 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 28 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Sections 1, 6 and 7.




Hole 361-U1479C Core 29H, Interval 257.5-267.43 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 29 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479C Core 30H, Interval 267.0-276.75 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 30 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 1.




Hole 361-U1479C Core 31H, Interval 276.5-286.49 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 31 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 1.




Hole 361-U1479C Core 32H, Interval 286.0-296.03 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 32 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 8/5GY; GLEY 1 8/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Change in lithification (from ooze to chalk) and color (from GLEY 1 8/5GY to GLEY 1 8/10Y) in Section 5 at 90 cm. Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 1 and 5-7.




Hole 361-U1479D Core 1H, Interval 0.0-9.39 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 1 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. One sandy interval in Section 2 at 75-84 cm. Moderate drilling disturbance in Section 1.



Hole 361-U1479D Core 2H, Interval 9.4-19.14 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 2 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate to extreme drilling disturbances in Sections 1-2 and 5-7.







Hole 361-U1479D Core 3H, Interval 15.9-24.68 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 3 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 1 and 5.











Hole 361-U1479E Core 2H, Interval 90.0-99.74 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 2 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 1 and 2.




Hole 361-U1479E Core 3H, Interval 99.5-109.07 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 3 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 1 to 3.




Hole 361-U1479E Core 4H, Interval 109.0-118.96 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 4 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 7 and CC.




Hole 361-U1479E Core 5H, Interval 118.5-128.47 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 5 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 6 to CC.




Hole 361-U1479E Core 6H, Interval 128.0-137.92 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 6 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 1 and 6 to CC.



Hole 361-U1479E Core 7H, Interval 137.5-147.48 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 7 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 1 to CC.






Hole 361-U1479E Core 8H, Interval 147.0-157.03 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 8 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 1 to CC.






Hole 361-U1479E Core 10H, Interval 158.5-168.2 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 10 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Sections 1 and 2.




Hole 361-U1479E Core 11H, Interval 168.0-178.02 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 11 comprises one lithological unit. Unit 1 is light greenish gray (GLE Y 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.



Hole 361-U1479E Core 12H, Interval 177.5-187.56 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 12 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Severe drilling disturbance in Section 1.




Hole 361-U1479E Core 133, Interval 187.0-229.0 m (CSF-A)

DRILLED INTERVAL

Depth CSF-A (m)	Core length (cm)	Section	Lithologic unit	Core image	Graphic lithology	Disturbance type	Disturbance intensity	Average grain size	Sedimentary structures	Lithological accessories	Bioturbation intensity	Shipboard samples	Reflectance			Magnetic susceptibility (SI)	GRA bulk density (g/m <sup>3</sup> )	
													L*	a*	b*			
221																		
222																		
223																		
224																		
225																		
226																		
227																		
228																		
229																		


Hole 361-U1479E Core 14H, Interval 229.0-238.94 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 14 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate to severe drilling disturbance in Sections 3 to 5.



Hole 361-U1479E Core 15H, Interval 238.5-248.52 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 15 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.






Hole 361-U1479E Core 16H, Interval 248.0-257.44 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 16 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate to extreme drilling disturbance in Section 1 and 3.




Hole 361-U1479E Core 17H, Interval 257.5-267.45 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 17 comprises one lithological unit. Unit 1 is light greenish gray (GLE Y 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.




Hole 361-U1479E Core 18H, Interval 267.0-276.87 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 18 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 6/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 7.




Hole 361-U1479F Core 1H, Interval 0.0-9.5 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 1 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. One sandy interval in Section 3 at 31-36 cm. Severe drilling disturbance in Section 1.




Hole 361-U1479G Core 1H, Interval 0.0-9.29 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 1 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. One sandy interval in Section 4 at 139-145 cm. Severe drilling disturbance in Section 1.



Hole 361-U1479H Core 1H, Interval 0.0-5.57 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 1 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. One sandy interval in Section 4 at 76-82 cm. Severe drilling disturbance in Section 1.





Hole 361-U1479H Core 3H, Interval 15.1-24.75 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 3 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.






Hole 361-U1479H Core 4H, Interval 24.6-34.24 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 4 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.



Hole 361-U1479H Core 5H, Interval 34.1-41.65 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 5 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Moderate drilling disturbance in Section 1 and severe in Section 5.






Hole 361-U1479H Core 7H, Interval 47.6-50.15 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 7 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Slight drilling disturbance in Section 1.




Hole 361-U1479H Core 8H, Interval 57.1-66.76 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 8 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.




Hole 361-U1479H Core 9H, Interval 66.6-73.09 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 9 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. One sandy interval in Section 5 at 22-28 cm. Extreme drilling disturbance in Section 4.




Hole 361-U1479H Core 10H, Interval 76.1-86.08 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 10 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.



Hole 361-U1479I Core 1H, Interval 0.0-6.31 m (CSF-A)


OOZE, NANNOFOSSILS, FORAMINIFERA Core 1 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 4.






Hole 361-U1479I Core 21, Interval 6.3-41.0 m (CSF-A)

DRILLED INTERVAL




Hole 361-U1479I Core 3H, Interval 41.0-43.35 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 3 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core.



Hole 361-U1479I Core 4H, Interval 50.5-60.03 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 4 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Sand intervals in Section 2 at 34-40 cm and in Section 3 at 30-33 cm. Extreme drilling disturbance in Section 1.



Hole 361-U1479I Core 5H, Interval 60.0-63.6 m (CSF-A)

OOZE, NANNOFOSSILS, FORAMINIFERA Core 5 comprises one lithological unit. Unit 1 is light greenish gray (GLEY 1 7/10Y) nannofossil ooze with foraminifera. Strong bioturbation is present throughout the Core (mainly burrows). Moderate diagenetic alterations (pyrite) occur at irregular intervals throughout the Core. Extreme drilling disturbance in Section 1.

