IODP-MSP (Exp. 364) VISUAL SECTION UNIT DESCRIPTION

Exp.	Site	Hole	Core	Type	Section
364	77	A	6	R	1

Date	Time	Observers
23/09/16		EC/MW

QUOTATION
COTE

[cm]	Image	Unit #	Lithology	Veins and Alteration	Structures	Burrows	Ichnofabrics	Fossils	Core Disturbance	Description
0–160		1	marlstone	11 and 21 cm veins / (calcite).						Laminated marlstone 2.5Y 3/2 very dark grayish brown. - Few nodules → don't react to HCl (≈ 2mm) - Between 22.5 and 25 cm, lighter interval 2.5Y 3/3 dark olive brown. CT facies: 0 to 10 cm = light grey; 10 to 22 cm = dark grey ⊕ a black line between 21–22 cm
		2	limestone 22.5cm – 25cm							coarsening up limestone (turbidite?) 2.5Y 5/2 grayish brown 25 to 24 cm: at the base, subangular clast (shell? other...) pyritised. ⊕ planar dark (organic?) 24 to 22.5: same observations. CT facies: white to light gray (high density high atomic n°)
		3	marlstone				1			Laminated marlstone with foram 2.5Y 3/2 ↳ several continuous and lenticular lighter beds: 2.5Y 6/2 (light brownish gray) (thickness 1 to 3 mm) at 41,5 cm, coarsening up (3mm thick). ↳ reaction to HCl - down up 51 cm. ↳ from 51 to 122,5 cm, laminated marlstone 2.5Y 3/2 with few burrow between (ichno 1) 66 cm et 77 cm → very dark grayish brown ⊕ few nodules (≈ 1 mm to 2 mm) ↳ don't reacted to HCl. CT facies: - white = 0 to 25 cm - light grey = 25 cm to 55 cm - dark grey to black = 55 cm to 70 cm - light grey = 70 cm to 120 cm
			122,5							GLEY 2 4/1
		4	claystone							sharp limitations at the base and the top → CT facies black; lighter interval 126/129 cm with pyrite (1 cm) → CT facies white; GLEY 2 5/6B 6
		5	129 marlstone							Marlstone laminated 2.5Y 3/2 very dark grayish brown with lighter interval 142 cm (1 cm thick) containing foram and clast (mm). 2.5Y 6/4 light yellowish brown.