IODP-MSP (Exp. 364) VISUAL SECTION UNIT DESCRIPTION

Exp.	Site	Hole	Core	Type	Section
364	77	A	23	R	2

Date	Time	Observers
25 09 16	14 35	EC MW



[cm] | Unit # | Lithology | Veins and Alteration | Structures | Burrows | Ichnofabrics | Fossils | Core Disturbance | Description

0 | 1 | marlstone | 22 30 cm | 2 to 3

Laminated marlstone bidurbated 4/2
- From 0 to 20 cm, fracture
CT facies black

20 | 2 | marlstone | 32/33 cm | 1 to 2

Laminated marlstone with local burrow (2.5Y 3/2)
- at 25 cm, claystone (1 cm-thick), uncontinuous.
CT facies blak

30 | 3 | marly packstone | marlstone 91 cm | 1 2 3 1

- Marlstone with very high carbonate content (2.5Y 7/2).
- At the base and at the top of the unit, we observed very planar laminated structure and darker (compression of burrows) (2.5Y 5/2)
(The whole unit is highly bidurbated)
CT Facies (high atomic n° + high density)
dark gray = limestone (lighter pal)
black = laminated base.

90 | 4 | marlstone | 103 to 110 cm | wavy base | 132 cm | 2 3 4 3 2

- Marlstone laminated bidurbated - (2.5Y 3/2)
From 103 to 110 cm, fracture
From 105 to 112 cm, lighter colod (2.5Y 5/2) by high content of carbonate.
- Claystone between 126 and 127 cm. (wavy)
CT facies.
dark gray claystone appear black

130 | 5 | marlstone | 1

Laminated dark marlstone with local burrows -
CT facies dark gray to black)

150

160