

































Hole 366-U1496A Core 11X, Interval 40.8-41.07 m (CSF-A)																		
greenish-gray serpentinite pebbly mud with lithic clasts																		
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance	Structures	Reflectance	Natural gamma radiation (cps)	GRA bulk density (g/cm³)	Magnetic susceptibility (SI)	Shear strength (kN/m²)			
							Dominant	Subdominant								Type	Intensity	
41	0	CC				IVC	10	5										


[illegible]

















Hole 366-U1496B Core 8X, Interval 27.4-27.64 m (CSF-A)																
Metabasalt with rodingite veins. Basalt greenstone is dark greenish gray (GLEY 2 4/10BG) medium grained rock crosscut by light green very thin veinlets and carbonate veins (up to 1 mm). It contains a rodingite domain. Rodingite composed of a reddish brown (10R 3/3 dusky red) mineral, green (Gley 1 3/5G very dark grayish green) chlorite or amphibole, and carbonate. Crosscut by carbonate veins.																
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance		Structures	Reflectance	Natural gamma radiation (cps)	GRA bulk density (g/cm³)	Magnetic susceptibility (SI)	Shear strength (kN/m²)
							Dominant	Subdominant	Type	Intensity		L* a* b*				
0	CC				IV				X	X						

Hole 366-U1496B Core 91, Interval 28.4-34.4 m (CSF-A)																		
DRILLED INTERVAL																		
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance	Structures	Reflectance			Natural gamma radiation (cps)	GRA bulk density (g/cm³)	Magnetic susceptibility (SI)	Shear strength (kN/m²)	
							Dominant	Subdominant			L*	a*	b*					
0																		
29																		
100																		
30																		
200																		
31																		
300																		
32																		
33																		
34																		

Fragments of ultramafic rock and metabasite (greenstone), presumed clasts in serpentinite mudflow. Ground up by rotary core bit.





Hole 366-U1496C Core 2R, Interval 8.0-17.5 m (CSF-A)																			
NO RECOVERY																			
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance	Structures	Reflectance			Natural gamma radiation (cps)	GRA bulk density		Magnetic susceptibility		Shear strength (kN/m²)
							Dominant	Subdominant			L*	a*	b*		MAD (g/cm³)	SHMSL (SI)			
8	0										0.0	0.0	0.0	0.00	1.00		100		0.0
9											0.3	0.3	0.3	0.25	1.50		1100		0.3
10											0.5	0.5	0.5	0.50	2.00		2100		0.5
11											0.8	0.8	0.8	0.75	2.50		3100		0.8
12											1.0	1.0	1.0	1.00	3.00		4100		1.0
13																			
14																			
15																			
16																			
17																			


Hole 366-U1496C Core 3R, Interval 17.5-27.1 m (CSF-A)																			
NO RECOVERY																			
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance	Structures	Reflectance			Natural gamma radiation (cps)	GRA bulk density (g/cm³)		Magnetic susceptibility (SI)		Shear strength (kN/m²)
							Dominant	Subdominant			L*	a*	b*		MAD	SHMSL			
0																			
18																			
19																			
20																			
21																			
22																			
23																			
24																			
25																			
26																			
27																			

Hole 366-U1496C Core 4R, Interval 27.1-36.6 m (CSF-A)																				
NO RECOVERY																				
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance		Structures	Reflectance			Natural gamma radiation (cps)	GRA bulk density		Magnetic susceptibility		Shear strength (kN/m²)
							Dominant	Subdominant	Type	Intensity		L*	a*	b*		MAD (g/cm³)	WRMSL (SI)	SHMSL		
0																				
28																				
29																				
30																				
31																				
32																				
33																				
34																				
35																				
36																				

Hole 366-U1496C Core 6R, Interval 46.1-55.6 m (CSF-A)																			
NO RECOVERY																			
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance	Structures	Reflectance			Natural gamma radiation (cps)	GRA bulk density		Magnetic susceptibility		Shear strength (kN/m²)
							Dominant	Subdominant			L*	a*	b*		MAD (g/cm³)		WRMSL (SI)	SHMSL (SI)	
	0										0.0	0.0	0.0	0.00	1.00		100		0.0
47											0.3	0.3	0.3	0.25	1.50		1100		0.3
48											0.5	0.5	0.5	0.50	2.00		2100		0.5
49											0.8	0.8	0.8	0.75	2.50		3100		0.8
50											1.0	1.0	1.0	1.00	3.00		4100		1.0
51																			
52																			
53																			
54																			
55																			

Hole 366-U1496C Core 7R, Interval 55.6-55.7 m (CSF-A)

Core U1496C-7R consists of a large clast of serpentinized harzburgite. It is coarse grained with a porphyroclastic texture, consisting of about 80% olivine/serpentine, 15% orthopyroxene/bastite and 5% spinel. Serpentinization is estimated to be 75%. Serpentine veins of numerous generations crosscut the clast.




Hole 366-U1496C Core 8R, Interval 65.3-74.8 m (CSF-A)																			
NO RECOVERY																			
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance	Structures	Reflectance			Natural gamma radiation (cps)	GRA bulk density		Magnetic susceptibility		Shear strength (kN/m²)
							Dominant	Subdominant			L*	a*	b*		MAD (g/cm³)	SHMSL (SI)			
	0																		
66																			
67																			
68																			
69																			
70																			
71																			
72																			
73																			
74																			


Hole 366-U1496C Core 9R, Interval 74.8-84.4 m (CSF-A)																		
NO RECOVERY																		
Depth (mbsf)	Core length (cm)		Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance	Structures	Reflectance			Natural gamma radiation (cps)	GRA bulk density		Magnetic susceptibility
								Dominant	Subdominant	Type	Intensity	L*	a*	b*		MAD (g/cm³)	SHMSL (SI)	Shear strength (kN/m²)
75	0																	
76																		
77																		
78																		
79																		
80																		
81																		
82																		
83																		
84																		

Hole 366-U1496C Core 10R, Interval 84.4-93.9 m (CSF-A)																			
NO RECOVERY																			
Depth (mbsf)	Core length (cm)	Section	Core image	Shipboard samples	Graphic lithology	Lith. unit	Clast >2mm abundance (%)		Drilling disturbance	Structures	Reflectance			Natural gamma radiation (cps)	GRA bulk density		Magnetic susceptibility		Shear strength (kN/m²)
							Dominant	Subdominant			L*	a*	b*		MAD (g/cm³)		WRMSL (SI)	SHMSL	
	0										0.0	0.0	0.0	0.00	1.00		100		0.0
85											0.3	0.3	0.3	0.25	1.50		1100		0.3
86											0.5	0.5	0.5	0.50	2.00		2100		0.5
87											0.8	0.8	0.8	0.75	2.50		3100		0.8
88											1.0	1.0	1.0	1.00	3.00		4100		1.0
89																			
90																			
91																			
92																			
93																			

Hole 366-U1496C Core 11R, Interval 93.9-99.66 m (CSF-A)

Core U1496C-11R consists of light greenish-gray serpentinite mud with about 2-5% clasts of serpentinitized ultramafic rock. Drilling disturbance is severe (core extension, void space and soupy sediments) throughout the core.





Core U1496C-13G consists of light bluish gray serpentinite mud with 2-3% granule to pebble size ultramafic rock clasts.

