Proceedings of the International Ocean Discovery Program, Volume 367/368

Expedition 367/368 Site U1504, Table T9. Distribution of calcareous nannofossils, Site U1504. Preservation: G = good. Abundance: A = abundant, C = common, F = few, R = rare, B = barren.

https://doi.org/10.14379/iodp.proc.367368.108.2018

| Hole, core, section | Top depth (m) | Bottom depth (m) | Preservation | Group Abundance | *Calcidiscus leptoporus* | *Calcidiscus premacintyrei* | *Ceratolithus cristatus* | *Clausicoccus subdistichus* | *Coccolithus miopelagicus* | *Coccolithus pelagicus* | *Coronocyclus nitescens* | *Cyclicargolithus abisectus* | *Cyclicargolithus floridanus* | *Discoaster berggrenii* | *Discoaster bollii* | *Discoaster brouweri* | *Discoaster deflandrei* | *Discoaster druggii* | *Discoaster exilis* | *Discoaster moorei* | *Discoaster pentaradiatus* | *Discoaster petaliformis* | *Discoaster prepentaradiatus* | *Discoaster quinqueramus* | *Discoaster sanmiguelensis* | *Discoaster signus* | *Discoaster* spp. | *Discoaster surculus* | *Discoaster variabilis* | *Emiliania huxleyi* | *Gephyrocapsa* sp. (<4 microns) | *Gephyrocapsa* sp. (>5.5 microns) | *Gephyrocapsa* sp. (4-5.5 microns) | *Helicosphaera ampliaperta* | *Helicosphaera carteri* | *Helicosphaera euphratis* | *Helicosphaera sellii* | *Orthorhabdus serratus* | *Pontosphaera multipora* | *Pontosphaera* spp. | *Pseudoemiliania lacunosa* | *Reticulofenestra asanoi* | *Reticulofenestra bisecta* (5-10 microns) | *Reticulofenestra lockeri* | *Reticulofenestra pseudoumbilicus* | *Reticulofenestra* sp. (5-7 microns) | *Reticulofenestra* spp. small | *Rhabdosphaera* spp. | *Scyphosphaera* spp. | *Sphenolithus abies* | *Sphenolithus belemnos* | *Sphenolithus disbelemnos* | *Sphenolithus heteromorphus* | *Sphenolithus moriformis* | *Sphenolithus pospichalii* | *Sphenolithus* spp. | *Syracosphaera* spp. | *Triquetrorhabdulus carinatus* | *Triquetrorhabdulus longus* | *Umbilicosphaera rotula* | *Umbilicosphaera sibogae* | Comments |
|---|---|---