

Core, section, interval (cm)	Top depth (m)	Bottom depth (m)	Preservation	Group abundance	Comments
368-U1505A-1H-CC, 0-5	0.33	0.38	G A	C	
368-U1505B-1H-CC, 6-22	3.18	3.23	G A	F C	
368-U1505C-1H-CC, 15-20	8.18	8.23	G A	F C	
2H-CC, 19-24	17.69	17.74	G A	F C	
3H-CC, 17-22	25.99	26.04	G A	F C	
4H-CC, 30-35	37.12	37.17	G A	F C	
5H-CC	46.46	46.51	G A	F C	
6H-CC	56.18	56.23	G A	F C	
7H-CC	65.65	65.70	G A	F C	
8H-CC	74.90	74.95	G A	F C	
9H-CC	84.13	84.18	G A	R F F	
10H-CC	93.60	93.65	G A	F C	
11H-CC	102.73	102.78	G A	F C	
12H-CC	111.68	111.73	G A	F C	
13H-CC	122.66	122.71	G A	F C	
14H-CC	132.18	132.23	G A	F C	
15H-CC	141.27	141.32	G A	F F	
16H-CC	150.85	150.90	G A	F F	
17H-CC, 34-39	160.62	160.67	G A	F F	
18H-CC, 29-34	170.23	170.28	G A	F F	
19H-CC, 33-38	179.70	179.75	G A	F F	
20H-CC, 30-35	189.21	189.26	G A	F F	
21H-CC, 17-22	196.48	196.53	G A	F F	
22H-CC, 25-30	200.01	200.06	G A	F F	
23H-CC, 37-42	207.00	207.05	G A	F F	
24F-CC, 24-34	212.13	212.18	G A	F F	
25F-CC, 32-37	216.87	216.92	G A	F F	
26F-CC, 24-29	221.49	221.54	G A	F F	
27F-CC, 33-38	226.28	226.33	G A	F F	
28F-CC, 22-27	230.88	230.93	G A	F F	
29F-CC, 27-32	235.64	235.69	G A	F F	
30F-CC	240.27	240.32	G A	F F	
31F-CC	244.97	245.02	G A	F F	
32F-CC	249.69	249.74	G A	F F	
33F-CC	254.41	254.46	G A	F F	
34F-CC	259.12	259.17	G A	F F	
35F-CC	263.78	263.83	G A	F F	
36F-CC	268.48	268.53	G A	F F	
37F-CC	272.96	273.01	G A	F F	
38F-CC	277.90	277.95	G A	F F	
39F-CC	279.35	279.40	G A	F F	
40F-CC	284.39	284.44	G A	F F	
41F-CC, 20-25	288.96	289.01	G A	F F	
42F-CC, 28-33	293.78	293.83	G A	F F	
43F-CC, 21-26	298.17	298.22	G A	F F	
44F-CC, 26-31	303.16	303.21	G A	F F	
45F-CC, 22-27	307.87	307.92	G A	F F	
46F-CC, 20-25	310.69	310.74	G A	F F	
47F-CC, 21-26	314.85	314.90	G A	F F	
48X-CC, 36-41	326.88	326.93	G A	F F	
49X-CC, 34-39	336.65	336.70	G A	F F	
50X-CC	346.07	346.12	G A	F F	
51X-CC	355.53	355.58	G A	F F	
52X-CC	364.93	364.98	M A	F F	
53X-CC	374.52	374.57	M A	F F	
54X-CC	383.9	383.95	M A	F F	
55X-CC, 44-49	394.06	394.11	M A	F F	
56X-CC, 37-42	403.79	403.84	M A	F F	
57X-1, 0-1	403.50	403.51	B		
57X-1, 24-25	403.74	403.75	B		
57X-1, 144-145	404.94	404.95	M R		
57X-6, 1-2	411.06	411.07	M C		
57X-6, 120-121	412.25	412.26	M C		
57X-7, 1-2	412.55	412.56	M C		
57X-CC, 1-2	413.06	413.07	M C		
57X-CC, 39-44	413.44	413.49	M A		
58X-CC, 37-42	419.53	419.58	M A		
59X-CC, 46-51	430.21	430.26	M A		
60X-CC	442.26	442.31	M A		
61X-CC	451.88	451.93	G A		
62X-CC	461.44	461.49	G A		
63X-CC	471.04	471.09	G A		
64X-CC, 36-41	480.49	480.54	G A		

Very good preservation of S. *Heteromorphus*
Very good preservation of S. *Heteromorphus*
Very good preservation of S. *Heteromorphus*

Overgrowth in discoasterids
Overgrowth in discoasterids
Overgrowths in discoasterids and *Sphenolithus*
Heavy overgrowth in discoasterids and *Sphenolithus*
Heavy overgrowth

Typical *Reticulofenestra isabellae*, *R. westerholdii*, *R. onusta*; *R. reticulata* not typical; *Cruciplacolithus tarquinius*, *C. klausii*, *Markalius inversus*