

Core, section, interval (cm)	Bottom depth CSF-A (m)	AW Planktonic (%)	OK Planktonic (%)	Planktonic mean (%)	Oceanicity	Paleowater depth (m)	
398-U1593A-							
1H-CC	4.73	NA	BARREN	NA	NA	NA	NA
2H-4, 63-65	9.86	88		88	Sub-oceanic		
2H-CC	13.30	50	46	48	Extra-neritic		
3H-CC	15.79	0	0	0	Inner neritic		
4H-CC	27.46	95	83	89	Sub-oceanic		
5H-CC	35.67	81	61	71			
6H-CC	42.83	15	12	13.5	Outer neritic		
7H-CC	62.09	65	60	62.5	Sub-oceanic		
8H-CC	70.01	27	17	22	Outer neritic		
9H-1, 60-62	71.82	60	38	49	Extra-neritic		
10H-CC	90.55	29	20	24.5	Outer neritic	NA	
11H-2, 35-37	92.07	15	24	19.5		NA	
11H-CC	94.66	58	BARREN	58	Extra-neritic	Reworked benthic	
12H-CC	109.18	57	43	50	Outer neritic	NA	
13H-2, 41-43	111.13	24	27	25.5		NA	
14H-CC	122.57	54	BARREN	54	Extra-neritic	Reworked benthic	
18F-CC	154.44	0	11	5.5	Inner neritic	NA	
25F-CC	187.87	31	BARREN	31	Extra-neritic	Reworked benthic	
26F-CC	194.18	29	38	33.5		NA	
28F-CC	203.39	0	33	16.5	Outer neritic	NA	
29F-CC	204.45	BARREN	BARREN	NA	NA	Reworked benthic	
30F-CC	210.46	17	BARREN	17	Outer neritic	NA	
31F-CC	215.47	28	25	26.5		Reworked benthic	
32F-CC	222.63	26	19	22.5		NA	
33F-CC	222.79	0	BARREN	0	Inner neritic	NA	
34F-CC	231.71	27	31	29	Outer neritic	NA	
35F-CC	235.54	32	38	35	Extra-neritic	Upper to lower bathyal	
36F-CC	241.50	81	80	80.5	Sub-oceanic	300->1000 m	
37F-CC	246.19	60	53	56.5	Extra-neritic		
38F-CC	250.89	85	60	72.5	Sub-oceanic		
398-U1593B-							
1H-CC	6.28		BARREN	NA		NA	NA
3H-CC	24.86		3	3.00		Reworked benthic	
7H-CC	63.70		14	14.00		NA	
9H-CC	82.34		39	39.00		NA	NA
10H-CC	92.08		24	24.00		Slightly dysoxic conditions	
11H-CC	101.72		42	42.00			
12F-CC	105.83		50	50.00			
13F-CC	109.62		59	59.00			
14F-CC	115.42		51	51.00			
25F-CC	165.58		19	19.00		NA	NA
26F-CC	167.28		19	19.00		Reworked benthic	
27F-CC	175.01		9	9.00			
28F-CC	178.67		10	10.00			
30F-CC	189.36		43	43.00			
33F-CC	202.07		BARREN	NA		NA	NA
34F-CC	208.47		BARREN	NA			
37F-CC	221.66		BARREN	NA			
38F-CC	226.74		20	20.00			
39F-CC	232.46		0	0.00		NA	NA

