








































Primarily silty clay with few sandy mud intervals and dispersed to common clasts throughout. There are small layers (~1-1.5 cm) containing coarser grains located at 15-16.5, 41-42, 60-61, 99-100.5, and 111-112 cm in Section 2, 171-72, 101-102 cm in Section 4, and very small (mm scale) authigenic iron oxides (?) visible on the surface of Sections 4, 5, 6, and 7. Sediments are structureless and firm with absent to moderate bioturbation and slight fracturing throughout.

































Hole 403-U1620A Core 24X, Interval 179.9-189.44 m (CSF-A)

Firm structureless silty clay with dark patches, some intervals with dispersed to abundant clasts. Color is mostly dark grey with small intervals of reddish sandy mud. Slight to heavy bioturbation. Moderate biscuiting and fracturing due to drilling occurs throughout all sections













Hole 403-U1620A Core 28X, Interval 208.9-218.72 m (CSF-A)


Primarily silty with with a ~50cm interval of sandy mud in Section 1 and muddy diamiction in Sections 2 and the top of 3. There is a 2-3cm sized sub-rounded clast at 52-55cm in Section 1 and a 3cm sized sub-rounded clast at 52-55cm in Section 3. Sediments are structureless and firm with no apparent bioturbation and strong biscuiting throughout.






































Hole 403-U1620C Core 8H, Interval 64.2-64.46 m (CSF-A)																
Recovery only includes core catcher. Soft structureless sandy mud of light grey color. No bioturbation.																
Depth CSF-A (m)	Core length (cm)	Section	Shipboard samples	Core image	X-radiograph image	Lithology	Disturbance type	Disturbance intensity	Bioturbation index	Sedimentary structures	Lower contact shape	Lower contact definition	Clast abundance	Magnetic susceptibility (IU)	NGR (cps)	Reflectance
	0		HS													

















Primarily silty clay with smaller intervals of sand and sandy mud. There is a 3cm sized sub-rounded clast at 68-70cm in Section 1, a 4cm sized sub rounded clast at 73-76cm in Section 2, and a 2cm sized sub rounded clast at 61-63cm in Section 3. Sediments are structureless other than faint laminations in Section 1, firm, and have no apparent bioturbation or drilling disturbance.




































Soft silty clay mostly with dark patches, some intervals with dispersed clasts. Thin layers of sand and sand patches in several sections. Color ranges from dark grey to dark yellowish grey with some greenish and reddish layers. No apparent to slight bioturbation.

























Firm dark grey structureless silty clay with thin intervals of reddish sand, dark patches, and dispersed to abundant clasts. Heavy to no bioturbation. Slight fracturing and biscuiting due to drilling occurs in some sections.































77


Silty clay with abundant authigenic iron oxide formation throughout. Sediments are firm and mostly structureless other than very faint, thin laminations in Section 6. There is absent bioturbation and strong biscuiting throughout.





















Firm dark grey structureless silty clay with dispersed to common clasts. Heavy to slight bioturbation. Strong biscuiting and fragmentation occur throughout all sections.























































































































Clayey silt with dispersed clasts in Section 1 and a gold colored 3-4cm size pyrite nodule cluster at 46-51cm. Sediments are hard and thinly laminated with absent to moderate bioturbation and biscuiting throughout.

