
































Primarily silty clay with dark patches and dispersed clasts with smaller intervals of sandy mud and one interval of interbedded silt (quartz) with silty clay in Section 2. Section 7 is biosiliceous-bearing. There is a 3 cm sized sub angular sand stone at 32-35cm in Section 3, small sand patches throughout Section 3, and a 5cm sub angular mud stone at 4-10cm in Section 7. Sediments are soft and primarily structureless with absent to slight bioturbation and up-arching throughout.




































Firm structureless light grey to dark grey silty clay with intervals of dark patches and dispersed clasts. There is little to no bioturbation. There is slight up arching and fracturing due drilling.

















Hole 403-U1621A Core 27X, Interval 195.9-196.6 m (CSF-A)													
Firm reddish sandy diamicton with no bioturbation or drilling disturbance. End of section 1 consists of only a hard rock likely plutonic.													
Depth CSF-A (m)	Core length (cm)	Section	Shipboard samples	Core image	X-radiograph image	Lithology	Disturbance type	Disturbance intensity	Bioturbation index	Sedimentary structures	Lower contact shape	Lower contact definition	Reflectance
									0 1 2 3 4				L* a* b*
196.00	0	1	HS MAD XRD CARB SED PAL NANNO FORAM DIAT										











Primarily silty clay with a smaller interval of clay in Section 4. There are dark patches throughout and dispersed clasts in a few sections. Similar to those found in Core 3, and Hole A, some smaller sections are composed of very black powdery sediments. There are small (~1-2 mm) irregular patches of coarser-grained sediment and the appearance of round white ~1-2mm in diameter authigenic minerals (?) apparent on surface of Section 6, and a large mafic 2.7cm clast at 42-46cm in Section 7. Sediments are structureless and soft with no apparent bioturbation and are slightly up-arched in some sections.





Silty clay with dark patches and dispersed clasts in most sections and interbedded silt in Section 1. There are jet black patches (iron sulfide?) in Section 1, a rounded 3cm clast (gray) at 120cm in Section 2, and a ~1.5cm rounded clast at 76cm in Section 4. Sediments are generally structureless and soft with no apparent bioturbation with slight to moderate up-arching and fracturing.


















































Intervals of silty clay, silt, clayey silt, and sandy mud with dispersed to common clasts throughout. There is a light brown layer from 32-33cm in the core catcher. Sediments are structureless and firm with absent to moderate bioturbation and slight fracturing and up-arching.






Intervals of silty clay and sandy mud with dispersed to abundant clasts throughout. There is a 1.5cm angular mudstone at 73-75cm in Section 2 and a 4 cm sized sub rounded sandstone at 84-88cm in Section 4. There are several thinly laminated intervals in Sections 1, 3, 4, and 6. Sediments are otherwise firm and structureless with absent to slight bioturbation, slight to moderate biscuiting, and slight up-arching and fall-in.








Intervals of silty clay, muddy diamicton, sandy mud, and clayey silt with dispersed to abundant clasts, shell fragments in the diamicton, a 3cm subangular gray clast at 46cm in Section 4, 2 2cm angular gray clasts from 82-84cm and a 4cm subangular gray clast at 130-134cm in Section 5. Sediments are structureless and firm with absent to heavy bioturbation and varying levels of fracturing throughout.
















Hole 403-U1621B Core 31X, Interval 201.1-202.24 m (CSF-A)													
Muddy diamicton with shell fragments throughout and a 2cm rounded granite at 10-11 and 2cm sub-angular mudstone at 69cm in Section 1 and a 2cm sized sub-rounded sandstone at 5-7cm and sub-angular 2cm sandstone at 13-14cm in the core catcher. Sediments are structureless and firm with no bioturbation and soupy sediments in the top 37cm of Section 1.													
Depth CSF-A (m)	Core length (cm)	Section	Shipboard samples	Core image	X-radiograph image	Lithology	Disturbance type	Disturbance intensity	Bioturbation index	Sedimentary structures	Lower contact shape	Lower contact definition	Clast abundance
									0 1 2 3 4				1 2 3
													Magnetic susceptibility (IU)
													19.6 21.6 23.6 25.6 39
													NGR (cps)
													44 49
													Reflectance
													L* a* b*
													-6.5 -4 -1.5
													28 33 38 43
													2 2.25 2.5 2.75 43
202.00	100	1	CC										


Washed cobble with no matrix. Composed of sandstone, mudstone, and quartzites.





Washed cobble with no matrix. Composed of sandstone, mudstone, and plutonic rocks.


















































Firm structureless grey silty clay to sandy mud with reddish intervals. Dark patches, dispersed clasts and sandy patches occur throughout all sections. A small interval of thin laminations occurs in section 4. There is no bioturbation. Slight fracturing a













Firm structureless grey silty clay to sandy mud with reddish intervals. Dark patches and sandy patches occur throughout all sections. A small interval of thin laminations occurs in section 3. There is little to no bioturbation. Slight fracturing and up ar

















This core only consists of gravel to cobble size rocks. The sediment matrix was washed away.

